

**Institut  
Français pour la  
Recherche et la  
Technologie  
Polaires**

**Centre des  
Faibles  
Radioactivités**

**Laboratoire mixte CNRS/CEA**

91198 Gif-sur-Yvette cédex

Les rapports de campagne à la mer  
à bord du *Marion-Dufresne*

**Campagne IMAGES  
MD 101**

du 29-05-95 au 11-07-95

**rédigé par  
Franck Bassinot et  
Laurent Labeyrie (chef de mission),  
avec les participants scientifiques.**

édition : juin 1996

Édité par l'Institut Français pour la Recherche et la Technologie Polaires  
Technopôle de Brest-Iroise - BP 75 - 29280 Plouzané - France  
Tél : (33) 02 98 05 65 00 - fax (33) 02 98 05 65 55



# **IMAGES MD 101**

**BREST-MARSEILLE  
29/05/95-11/07/95**

## **A CORING CRUISE OF the R/U MARION DUFRESNE in the NORTH ATLANTIC OCEAN AND NORWEGIAN SEA**


Cruise report prepared by F. Bassinot<sup>1</sup>, L. Labeyrie<sup>1,2</sup>,  
and the shipboard scientific party<sup>3</sup>

<sup>1</sup> Centre des Faibles Radioactivités, Laboratoire mixte CNRS-CEA,  
91198 Gif/Yvette cedex, France

<sup>2</sup> also at Dept. des Sciences de la Terre, Bat. 504, Université Paris Sud-  
Orsay, 91405 Orsay cedex, France

<sup>3</sup> see names and adresses in pages 4 and 5 of this report

Many thanks go to C. Bassinot, B. Lehman and J. Tessier from the  
CFR for their help in the final stage of preparation of this report



# TABLE OF CONTENTS

## SOM MAIRE

<b>* LIST OF SHIPBOARD SCIENTISTS</b>	4
<b>* CRUISE REPORT (english version)</b>	
1. Objectives of the IMAGES program	7
2. The IMAGES-MD101 cruise: objectives and results	7
3. Areas studied during IMAGES -MD101 cruise	8
<b>* RAPPORT DE CAMPAGNE (version française)</b>	
A. Preparation de la campagne MD101	11
1. Bases scientifiques: le programme IMAGES	11
2. Bases stratégiques	12
3. Bases financières et organisation pratique	12
B. Programme de recherche et déroulement de la campagne	12
1. Objectifs scientifiques principaux	12
2. Stratégie d'approche et méthodes	13
3. Objectifs détaillés, zone par zone	14
<b>* STATION AND CORE TABLES</b>	17
A. Summary table of journeys, stations and operations	17
B. Summary table of stations (operation, problems, ...etc)	23
C. Table of giant piston cores (Calypso cores)	28
D. Table of gravity cores	29
E. Table of plankton tows (and core catcher samples)	29
<b>* ON BOARD MEASUREMENTS</b>	
A. Preliminary report on physical properties and color imaging	31
<i>(G. Dunbar, F. Bassinot, K. Jarrett and the shipboard scientific party)</i>	
1. Continuous measurements of physical properties (MST)	31
1.1 Description of the GEOTEK multi-sensor track	31
1.2 Core handling	33
1.3 Calibration and initial setting of the MST	33
1.4 MST logging procedure	33
1.5 Data processing	34
1.6 Problems	35
2. Discrete measurements of index properties	36
3. Spectrophotometry	36
4. Video imaging and pictures	36
5. References	37
B. Preliminary report on calcareous nannofossil biostratigraphy	39
<i>(L. Beaufort and J. Giraudeau)</i>	
Biostratigraphy summary table:	40
<b>* SITE REPORTS</b>	43
A. Site reports format	43
B. Sedimentological charts	44
<i>(Cremer, M. and the shipboard sedimentological scientists)</i>	
C. Core locations	45
D. Site reports	46 to 217

## Shipboard participants to IMAGES-MD101 cruise

Participants MD101	Addresses	e-mails
Fatima Abrantes	Serv. Geol. Portugal, Nucleo de Geol. Marhina, Rua da Aca. das Ciencias, 19-2, P-1200 Lisboa, Portugal	
Jeff Adkins	Dept. of Earth, Atmospheric and Planetary Sciences, MIT, Cambridge, MA 02139, USA	adkins@mit.edu
Carin Andersson	Dept. of Geology, Univ. of Bergen, Alligaten 41, N-5007 Bergen, Norway	Carin.Andersson@tele.su.se
Harald Andruleit	SFB 313, Univ. Kiel, Olshausenstrasse 40, Kiel 23000, Germany	handruleit@geomar.de
G�rard Auffret	IFREMERM, Lab. Environnements s�dimentaires, Dept. Geosci. Marines, BP 70, 29280 Plouzan�, Fr	gauffret@ifremer.fr
Bill Austin	Dept. Geology and Geophysics, Univ. Edinburgh, Grant Institut, West Main Road, Edinburgh EH9 3JW, UK	Wena@glg.ed.ac.uk
Jeff Berry	Dept. of Earth, Atmospheric and Planetary Sciences, MIT, Cambridge, MA 02139, USA	jberry@mit.edu
Jacob Baas	GEOMAR, Wischhofstrasse 1-3, D-24148 Kiel, RFA	jbaas@geomar.de
Franck Bassinot	Centre des Faibles Radioactivit�s, Domaine du CNRS, 91198 Gif-sur-Yvette, France	bassinot@cfr.cnrs-gif.fr
Luc Beaufort	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	beaufort@arbois.cerege.fr
Guy Bilodeau	GEOTOP, Univ. Quebec, CP 8888 Success Avenue, Montreal, PQ H3C 3P8, Canada	r21360@er.uqam.ca
An Boelaert	IFREMERM, Lab. Environnements s�dimentaires, Dept. Geosci. Marines, BP 70, 29280 Plouzan�, Fr	An.Boelaert@ifremer.fr
Ed Boyle	Dept. of Earth, Atmospheric and Planetary Sciences, MIT, Cambridge, MA 02139, USA	eaboyle@mit.edu
No�lle Buchet	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	nbuchet@arbois.cerege.fr
Olivia Cayre	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	ocayre@arbois.cerege.fr
Herv� Cambray	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	cambray@imtmerl.imt-mrs.fr
Sylvie Chassery	UA 197 du CNRS, Dept. Geol. et Oceanologie, Av. des Facult�s, Univ. Bordeaux 1, 33405 Talence, Fr	(grousset@bxnms.u-bordeaux.fr)
Ester Clavero	Dept. Quimica Ambiental, CID-CSIC, Jordi Girona, 18. 08034-Barcelona, Spain	?
Michel Cremer	UA 197 du CNRS, Dept. Geol. et Oceanologie, Av. des Facult�s, Univ. Bordeaux 1, 33405 Talence, Fr	cremer@geocean.u-bordeaux.fr
Michel Decobert	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	mdecober@arbois.cerege.fr
Bernard Denielou	IFREMERM, Lab. Environnements s�dimentaires, Dept. Geosci. Marines, BP 70, 29280 Plouzan�, Fr	denielou@ifremer.fr
Shirley Derrick	Dept. Geology and Geophysics, Univ. Edinburgh, Grant Institut, West Main Road, Edinburgh EH9 3JW, UK	?
Derek Dregger	Geologisch-Paleontologisches Institut., Olshausenstrasse 40, Kiel 23000, Germany	dregger@sfb313.uni-kiel.d400.de
Gavin Dunbar	Dept. of Geology, James Cook Univ., Townsville, Q.4811, Australia	Gavin.Dunbar@jcu.edu.au
Mary Eliot	Centre des Faibles Radioactivit�s, Domaine du CNRS, 91198 Gif-sur-Yvette, France	Eliot@cfr.cnrs-gif.fr
Juan-Abel Flores	Depart. de Geologia, Fac. Ciencias, Univ. Salamanca, 37008-Salamanca, Spain	flores@rs6000.usal.es
Gerard Foch	IFREMERM, Lab. Environnements s�dimentaires, Dept. Geosci. Marines, BP 70, 29280 Plouzan�, Fr	?
Torben Fronval	Dept. of Geology, Univ. of Bergen, Alligaten 41, N-5007 Bergen, Norway	torben.fronval@geol.uib.no
Jacques Giraudeau	UA 197 du CNRS, Dept. Geol. et Oceanologie, Av. des Facult�s, Univ. Bordeaux 1, 33405 Talence, Fr	giraudeau@geocean.u-bordeaux.fr
Jens Gr�tzn�r	GEOMAR, Wischhofstrasse 1-3, D-24148 Kiel, RFA	jgruetzner@geomar.de
Heidrun Hensch	Geologisch-Paleontologisches Institut., Olshausenstrasse 40, Kiel 23000, Germany	?
Reinhard Hesse	Earth & Planet. Sciences, McGill Univ., 3450 Montreal, Quebec, H3A 2A7, Canada	Rein_H@geosci.lan.McGill.ca
Kate Jarrett	Atlantic geoscience Centre, Bedford Inst. Oceanography, PO Box 1006, Dartmouth, N.S. B2Y 4A7, Canada	jarrett@agcux.bio.ns.ca
Saeed Khodabakhsh	Earth & Planet. Sciences, McGill Univ., 3450 Montreal, Quebec, H3A 2A7, Canada	?
Nolan Koc	Dept. of Geology, Univ. of Bergen, Alligaten 41, N-5007 Bergen, Norway	Nalan.Koc@geol.uib.no

Participant MD101	Address	e-mails
Boris Konyukhov	Earth & Planet. Sciences, McGill Univ., 3450 Montreal, Quebec, H3A 2A7, Canada	?
Laurent Labeyrie	Centre des Faibles Radioactivités, Domaine du CNRS, 91198 Gif-sur-Yvette, France	labeyrie@cfr.cnrs-gif.fr
Yves Lancelot	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	ylancelo@arbois.cerege.fr
Irène Lefevre	Centre des Faibles Radioactivités, Domaine du CNRS, 91198 Gif-sur-Yvette, France	Lefevre@cfr.cnrs-gif.fr
Flavien Lemoine	Centre des Faibles Radioactivités, Domaine du CNRS, 91198 Gif-sur-Yvette, France	lemoine@cfr.cnrs-gif.fr
Linda Levesque	GEOTOP, Univ. Quebec, CP 8888 Success Avenue, Montreal, PQ H3C 3P8, Canada	m107261@er.uqam.ca
Ph. Martinez	UA 197 du CNRS, Dept. Geol. et Oceanologie, Av. des Facultés, Univ. Bordeaux 1, 33405 Talence, Fr	(bertrand@bxnms.u-bordeaux.fr)
Larry Mayer	Ocean Mapping Group, Univ. New Brunswick, Fredericton, N.B., E3B 5A3, Canada	lmayer@jupiter.sun.csd.unb.ca
Jurgen Mienert	GEOMAR, Wischhofstrasse 1-3, D-24148 Kiel, RFA	jmiert@geomar.de
Stefan Nees	GEOMAR, Wischhofstrasse 1-3, D-24148 Kiel, RFA	Snees@geomar.de
Nicole Page	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	npage@arbois.cerege.fr
Martine Paterne	Centre des Faibles Radioactivités, Domaine du CNRS, 91198 Gif-sur-Yvette, France	Paterne@cfr.cnrs-gif.fr
Carles Pelejero	Dept. Quimica Ambiental, CID-CSIC, Jordi Girona, 18. 08034-Barcelona, Spain	?
Severine Pennon	Centre des Faibles Radioactivités, Domaine du CNRS, 91198 Gif-sur-Yvette, France	?
Uwe Pflaumann	Geologisch-Paleontologisches Institut., Olshausenstrasse 40, Kiel 23000, Germany	up@zaphod.gpi.uni-kiel.de
Frank Rack	Ocean Mapping Group, Univ. New Brunswick, Fredericton, N.B., E3B 5A3, Canada	rack@omg.unb.ca
(Marie Revel)	UA 197 du CNRS, Dept. Geol. et Oceanologie, Av. des Facultés, Univ. Bordeaux 1, 33405 Talence, Fr	(grousset@bxnms.u-bordeaux.fr)
Michael Sarnthein	Geologisch-Paleontologisches Institut., Olshausenstrasse 40, Kiel 23000, Germany	MSarnthein@zaphod.gpi.uni-kiel.de
Nick Shackleton	Godwin laboratory, Free school line, Cambridge, Eng CB2 3RS, UK	NJS5@cam.ac.uk
Joachim Schönfeld	GEOMAR, Wischhofstrasse 1-3, D-24148 Kiel, RFA	jschoenfeld@geomar.de
Francisco Sierro	Depart. de Geologia, Fac. Ciencias, Univ. Salamanca, 37008-Salamanca, Spain	Sierro@rs6000.usal.es
Peter Solheid	Inst. Rock Magnetism, 293 Shepherd Labo., 100 Union Street S.E., Minneapolis, MN 55455-0128, USA	sohle003@maroon.tc.umn.edu
Colin Summerhayes	Southampt. Oceanog.Cent. (SOC), Empress Dock, Southampton, SO14 3ZH, United Kingdom	Colin.P.Summerhayes@soc.soton.ac.uk
Nicolas Thouveny	Labo. Geol. du Quaternaire, CEREGE, BP 80, 13762 Les Milles cedex, France	nthouven@arbois.cerege.fr
Jean-Louis Turon	UA 197 du CNRS, Dept. Geol. et Oceanologie, Av. des Facultés, Univ. Bordeaux 1, 33405 Talence, Fr	turon@geocean.u-bordeaux.fr
Mara Weinelt	Geologisch-Paleontologisches Institut., Olshausenstrasse 40, Kiel 23000, Germany	?
Claudia Willamowski	GEOMAR, Wischhofstrasse 1-3, D-24148 Kiel, RFA	?



## CRUISE REPORT (english version)

### THE FIRST IMAGES GIANT CORING CRUISE ON BOARD R/V MARION DUFRESNE (MD 101): HIGH RESOLUTION PALEOCEANOGRAPHY OF THE NORTH ATLANTIC .

**Labeyrie, L.** Centre des Faibles Radioactivités (CFR), CNRS/CEA, 91198 Gif/Yvette cedex and Département des Sciences de la Terre, Université Paris sud-Orsay, 91405 Orsay, France, Chief Scientist; **J.L. Turon** and **Y. Lancelot**, co-chief scientists; **F. Bassinot**, assistant Chief Scientist, **Y. Balut** Director for oceanographic operations and the **scientific party of IMAGES MD 101**.

#### 1/ IMAGES objectives.

The IMAGES program addresses two major questions for paleoceanography:

- How have changes in ocean properties controlled the evolution of global heat transfer through the deep and surface ocean and so modified climate.
- How have changes in ocean circulation, ocean chemistry, and biological activity interacted to generate the observed record of atmospheric pCO<sub>2</sub> over the past 300 ka?

IMAGES aims at a global integration of ocean sediment records with the Greenland and Antarctic ice records within the climate dynamic modeling effort. A temporal resolution of the order of centuries or better is a prerequisite for these studies. IMAGES recommends as a strategy the continuous sampling and analysis of high resolution records with better than 10 cm/kyr sedimentation rates. Records longer than 250 kyr are needed to differentiate the linear response to the insolation external forcing from the non linear interactions of ice, ocean, atmosphere and continental crust systems. This demands long sediment cores (several tens of meters) and a combination of precise geochemical and micro-paleontological methods applied on sets of samples at discrete intervals (2 to 5 cm). Continuous and non-destructive physical properties records obtained on board on sediment cores will serve as proxies for high resolution (cm) interpolation of geochemical and micro-paleontological results.

#### 2/ The IMAGES-MD101 cruise : objectives and results.

IMAGES coordinated a first international cruise in June and July 1995 over the North Atlantic and Norwegian sea on board the French R/V *Marion Dufresne* (MD 101, Brest - Stornoway (Lewis Island) - St-Pierre - Azores - Marseille).

Its main scientific objective was the collection of giant piston cores on rapidly sedimenting drifts and continental margins of the North Atlantic ocean and Norwegian Sea, along the track of the main thermohaline circulation. The cruise crossed the North-East Atlantic margins, the Feni Drift, the Scottish, North Faeroes and Norwegian margins (to 72°N), the Iceland South-East margins, the Gardar Drift, the NAMOC Channel, the Newfoundland margin, the Bermuda rise, the mid Atlantic ridge, and the Azores and Iberian margins.

Additional objectives covered :

- the contribution of Mediterranean waters to the North Atlantic intermediate waters, with 5 cores recovered across the slopes of the Iberian margin;
- the evolution of the NAMOC channel, in the deep North-West Atlantic basin, in relation to the growths and decays of the Laurentide ice sheet (8 Kullenberg and gravity cores).

This was the maiden cruise of the new *Marion Dufresne*, just 2 weeks out from her Le Havre shipyard. The ship had a very small number of problems, taking into account the number of things which were not ready just a few days before the departure. Two days were lost for engine problems.

70 scientists, students and technicians from 22 institutions (13 countries) participated to at least one of the three legs. 43 cores (mean length over 30 meters) have been retrieved during the cruise, described and measured for magnetic susceptibility, p-wave velocity,  $\gamma$  density and spectral light reflectance. The longest core, MD 95-2036 (52.64 m) was retrieved at 4461 m water depth on the Bermuda Rise. It covers about 150 kyr with a sedimentation rate over 30 cm/kyr. The Calypso corer worked properly, once a few problems encountered at the beginning of the cruise had been solved (i.e. sliced or imploded PVC liner).

This report presents preliminary results, mostly obtained on board: core descriptions, physical properties and micro-paleontological stratigraphy.

Color reflectance (between 40° and 55°N) and magnetic susceptibility (between 50° and 70°N) have been used for direct tuning of the time scales by cyclo-stratigraphy in the precession and obliquity bands. Ocean-wide correlations have been established over the last 250 kyr.

### **3/ Areas studied during the IMAGES-MD101 cruise.**

**Area 1 : Celtic continental slope at the southern end of the Channel (Trevelyan area, 3788 m, Core MD95-2001; Meriadzec area, 2174 m, Core MD95-2002).**

These sites which present exceptionally high sedimentation rates (on the order of 50 cm/kyr during isotopic stage 3) made it possible to test the Calypso giant coring system. They are particularly suitable for high resolution studies of climatic and sedimentary variability during the last glacial period.

**Area 2 : Feni Drift, east of Rockall Plateau (55° N 15°W ):**

This area is located along the initial pathway of the NADW in its journey towards the South and the West, and along the pathway of the main warm surface current of the North Atlantic (the North Atlantic Drift) in its journey towards the Norwegian Sea. Numerous cores from about 5 to 10 m in length have been studied previously on this sediment drift which is characterized by high sedimentation rates but irregular sedimentation processes. This area is a top priority for the CFR-DGO and German teams. About 5 cores were initially planned but, because of the 2 days of delay at the beginning of the cruise, only 2 cores have been retrieved (MD 95-2003 et 2004).

**Area 3 : north -west of Scotland:**

Coring of near coastal areas with very high sedimentation rates for the detailed study (temporal resolution on the order of 100 years) of climatic evolution since the last glacial maximum. Barra Fan, : Lat. 57°018N, 10°035 W, 2000 m (MD 95-2005 -boat drifted out of the planned target site- and MD 95-2006) and St Kilda Basin: 57°31'N, 08°23'W 158m (MD 95-2007). Objectives of the University of Edinburgh (W. Austin).

**Area 4 : Faeroes Ridge:**

This area has already been studied based on short piston cores. This sill located between the North Atlantic and the Nordic seas has an exceptional interest because the its sediments have recorded both variations of the northward penetration of surface waters from the north Atlantic and the response of deep waters. Two cores retrieved at about 62°58N, 2°22W, 1760m (MD 95-2008 et MD 95-2009). Sedimentation rate on the order of 10 cm/kyr. Top priority for the CFR-DGO group and the German teams (GEOMAR, Univ. of Kiel).

**Area 5 : Vöring Plateau.** This area is already well known from the study of short piston cores and from ODP Site 644. It presents an exceptional interest for the study of the evolution of the Scandinavian ice sheet in relationships with Greenland ice records (GISP and GRIP) and the changes of the thermohaline circulation in the Norwegian Sea. The sedimentation rates are high. Two sites were cored: MD 95-2010 66°58N, 07°38 E, 1048 m ; MD 95-2011 66°41N, 04°34 E, 1226 m. Priority of the Univ. of Bergen (E. Jansen) and the German laboratories (GEOMAR, Univ. of Kiel).

**Area 6 : Bear Island.** This area, which was located in front of the small Barents Sea ice cap during the last glacial period, plays an important role in the processes linked to the early deglaciation. The coring of a site with a high sedimentation rate was the main objective of the team from the Kiel University. Unfortunately, the two cores retrieved were badly damaged (PVC liners were badly sliced and imploded). (MD 95-2012 72°09N, 11°26E, 2094m et MD 95-2013 72°11N, 13°06E, 1638 m).

**Area 7 : southern margin of Iceland (Reykjanes, 60°N-22°W).** This area -which shows sedimentation rates on the order of 8-12 cm/kyr- has been previously studied during the Paleocinat 1 cruise of the R/V Suroît (cores SU 90-32 and 33, 10 and 15 m in length, respectively). This area makes it possible to study, over a few climatic cycles, the northward extension of the warm surface waters of the North Atlantic Drift, the links with the evolution of the Iceland ice sheet, and the intermediate waters history. Numerous ash layers can be dated with the Ar<sup>39</sup>-Ar<sup>40</sup> method to improve the time scale derived from orbital tuning of the isotopic stratigraphy. This area is a top priority for the CFR-DGO group. One core has been retrieved on the same location as core SU 90-33 (MD 95-2014 60°35 N, 22°05W, 2397 m).

**Area 8 : Gardar Drift and mid-Atlantic Ridge (55°-57°N : 30-40°W).** The last results obtained by the CFR-DGO group tend to show that deep water formation took place in this area during glacial periods. The mechanism is close to the one observed today in the Labrador Sea (modal waters). There was no good core available in this area prior to the IMAGES-MD101 cruise. This is a top priority objective for the CFR-DGO group. Two cores have been retrieved: MD95-

2015 à 58°46 N 25°58 W 2630 m et MD95-2016 à 57°43 N et 29°26 W 2318 m. The latter one has been retrieved on a location already studied by W. Ruddiman. The sedimentation rate is exceptionally high during interglacials (higher than 30 cm/kyr).

Areas 9-10 : north-east margin of North America (44°-50°N : 40°-55°W). Sediments of these areas provide the best records of the massive iceberg discharges which corresponded to the "Heinrich events" initiated in the northern part of the Laurentide. At deep water depths, one can analyze the variations of the NADW in its southward track, out of its formation areas. The sites have been selected based on previous records from numerous cores obtained by the Lamont (LDEO, Univ. of Columbia, NY), the Canadian teams, or by the CFR-DGO during the Paleocinat 1 cruise. This area represents the unique objective of the Canadian teams involved in the IMAGES program (Univ. Of Fredericton, Univ. of New Brunswick, GEOTOP). A "non-IMAGES" objective was raised by a group from McGill University (Can.): a detailed sampling of the NAMOC canyon. This canyon follows on more than 2000 km the slope between the Labrador margin and the abyssal plain of the western Atlantic basin. It has been apparently dug by massive arrivals of turbid waters that flew out of the base of the Laurentide ice sheet during its periods of maximal extension. 8 cores have been retrieved on the NAMOC zone (MD 95-2018 à 23). The first core retrieved for the Canadian IMAGES program was located at one edge of the Gibbs fracture zone, on the mid-Atlantic Ridge, south-east of Gardar Drift (MD 95-2017 53° 03 N, 33°32W, 3100 m). Other cores were retrieved on the Canadian margin, north, east and south-west of Newfoundland, between 4150 and 878 m of water depth (cores MD 95 2024 to 2033).

Area 11 : Bermuda Rise. This high sedimentation rate area (> 30 cm/kyr) permits to reconstruct with a very good resolution the variations of surface subtropical waters from the Gulf Stream and the variations at depth of the boundary between the NADW and the AABW. Three cores have been obtained in order to cover the last climatic cycles (MD95-2034 to MD95-2036). The Core MD95-2036 is 52.64 m in length, the record for Kullenberg coring in the North Atlantic. These sites were the main objectives of the US teams.

Areas 12-13: Mid-Atlantic Ridge, Azores and western Iberian margin. The Azores and the Iberian margin areas have been previously studied during the 1980 and 1990 cruises of the R/V Suroît and cruises of the R/V Meteor. They present high sedimentation rates (between 10 and 50 cm/kyr). These areas are ideally located to study the evolution of the southernmost limit of the Polar Front during glacial periods and the contribution of the Mediterranean waters to the intermediate waters of the Atlantic. The sediments from the Azores have an exceptional interest for paleovolcanic and paleomagnetic studies. These areas are top priorities for the LGQ, the URA Orsay-Muséum, Ifremer-Brest, LODYC-Paris, the CFR-DGO group, the German (GEOMAR) and Portuguese teams (Portugal Geol. Surv.), and the British teams from the IOS and the Univ. of Cambridge. Between 8 and 10 cores were originally planned in this area, but only 6 could be retrieved because of the delay accumulated during the cruise (MD 95-2037 - 2042).

Area 14 : Western Mediterranean Sea. A high accumulation rate area in which it will be possible to reconstruct the paleoclimatic evolution of the southern part of Europe (based on pollens and bio-markers) in relationship with the Mediterranean hydrology. Priority of the Universities of Salamanca and Barcelona (and collaboration with Brest-Ifremer) : MD 95-2043 à 36°08 N, 02°37W, 1841 m).



# RAPPORT DE CAMPAGNE (version française)

## A - PREPARATION DE LA CAMPAGNE IMAGES-MD101

### 1/ Bases scientifiques: le programme IMAGES:

La communauté internationale des chercheurs travaillant sur les changements climatiques est de plus en plus concernée par le manque actuel de connaissance sur les origines, les mécanismes, et les conséquences des changements climatiques rapides observés dans les enregistrements paléoclimatiques (glaces, océans et continents). Il ne fait plus de doute que l'océan joue un rôle majeur dans ces changements, tant pour les transferts de chaleur que pour les contraintes physico-chimiques à long terme sur la teneur en CO<sub>2</sub> atmosphérique. Mais ce rôle est très imparfaitement évalué, faute de données suffisantes sur la variabilité dynamique et physico-chimique des océans. Les derniers résultats publiés dans Nature (Bond et al 1993, Grip 1993..) montrent l'importance probable du couplage entre la formation des eaux profondes nord atlantiques (North Atlantic Deep Water, NADW) et la dynamique des calottes de glace de la Laurentide, du Groenland et de la Fenno-Scandie; mais les transferts de chaleur entre hémisphères nord et sud, les flux de chaleur sensible et latente vers l'atmosphère, l'accumulation de CO<sub>2</sub> dissous dans les océans profonds, sont d'autres facteurs à considérer.

Les outils pour étudier les systèmes passés existent : composition isotopique de l'oxygène et du carbone des foraminifères planctoniques et benthiques, micropaléontologie, biomarqueurs (Uk37) pour reconstruire les gradients de température et salinité, les flux de carbone, et la paléoproduktivité, traceurs magnétiques, minéralogiques et géochimiques pour reconstruire les apports des continents.

Toutefois les carottes de sédiment existant dans les collections ne suffisent pas à ces études : les séries sont souvent incomplètes, et pas assez de matériel n'est disponible pour réaliser à très haute résolution les analyses nécessaires. De plus, ces carottes sont pour la plupart trop courtes pour couvrir une période de temps suffisante (au moins les deux derniers cycles glaciaires-interglaciaires) avec la résolution temporelle nécessaire à l'étude des changements du climat : cent à deux cent ans.

Ces réflexions ont conduit à la création du programme international IMAGES (International marine Global Change Study), animé par IGBP-PAGES (Past Global Changes) et SCOR (Scientific Committee on Oceanic research) et du programme US-NSF MESH (Marine Earth System History). Ces programmes mettent en priorité l'acquisition d'un grand nombre de carottes de sédiment marin (2000 pour les prochains 5-10 ans) de grande longueur, et disponibles pour toute une série d'analyses à très haute résolution permettant de décrire la variabilité des océans au cours des derniers 300 000 ans.

En France, le programme paléoclimatologie du PNEDC a lui aussi mis en toute première priorité la description de la variabilité hydrologique des masses d'eau superficielles et profondes de hautes latitudes, en particulier en Atlantique nord. Le Comité Scientifique International du PNEDC recommande une participation forte dans le programme de carottage IMAGES. La campagne Paléocinat, en cours d'exploitation, a fourni des informations très importantes, mais les carottes obtenues sur le SUROIT sont trop courtes (10 à 15 mètres): les derniers interglaciaires n'ont pu être échantillonnés que dans les zones à faibles vitesses de sédimentation, où la résolution est insuffisante pour des études paléoclimatiques quantitatives. Des carottages de grande longueur (40 m ou plus) sont indispensables à la poursuite du projet.

## **2/ Bases stratégiques:**

La proposition IMAGES-MD 101 a été le résultat de plusieurs facteurs:

a/- Le Marion Dufresne II était disponible à partir de la France (Brest) au printemps 1995. Il devait disposer d'un sondeur multi-faisceaux moderne et de possibilités de carottages encore améliorées.

b/- IGBP-PAGES (Past Global Change) et SCOR ont décidé de lancer en septembre-octobre 1993 le programme international IMAGES d'étude du rôle de l'océan dans les changements climatiques. Le groupe de travail WG 100 du SCOR, qui en était le comité de planification, a mis comme première priorité la réalisation de campagnes internationales de carottage géant et d'étude à très haute résolution des sédiments le long de 5 profils à travers les différents océans. Il a été proposé d'obtenir de différentes agences scientifiques, au niveau international, un support financier suffisant pour permettre de réaliser la première campagne sur le MD, seul navire au monde pouvant réaliser en routine des carottes de 50 mètres. Il s'agissait de réaliser en priorité une couverture des différentes zones de l'Atlantique nord. La campagne devait servir de plate-forme de lancement pour le programme IMAGES, avec essais simultanés des différents modes de prélèvement et d'étude en continu des carottes de sédiment. La Grande Bretagne (NERC), les USA (NSF), le Canada (NSERC), l'Allemagne, l'Espagne et la Norvège ont donné leur accord pour participer à un financement international du MD.

c/ En France, la campagne a été approuvée par l'IFRTP et le PNEDC (C.S. décembre 1993), après examen très favorable de la Commission spécialisée Géosciences Marines de l'IFREMER.

## **3/ Bases financières et organisation pratique:**

Les règles d'organisation et de financement de cette campagne "test" n'ont pas été celles qui seront adoptées définitivement pour les campagnes IMAGES à venir (encore en cours de discussion). A l'étape préliminaire actuelle, chaque investigateur principal étranger a financé par ses supports nationaux une quote-part, à partir d'un projet de participation sur des objectifs précis. Toutefois, les investigateurs principaux sont ensemble co-responsables des opérations de type IMAGES, en particulier l'étude en continu des paramètres physiques du sédiment et la stratigraphie, et des collaborations sont engagées sur l'ensemble du matériel prélevé. La faisabilité de ce type de collaboration est un test majeur pour la faisabilité du programme IMAGES.

## **B - PROGRAMME DE RECHERCHE ET DEROULEMENT DE LA CAMPAGNE**

### **1/ Objectifs scientifiques principaux:**

a-Reconstruction de la variabilité de l'hydrologie de surface de l'Atlantique nord, réponse de la circulation profonde, et évolution simultanée de la climatologie péri-Atlantique nord au cours des derniers cycles climatiques (objectif IMAGES).

Le thème principal de la campagne MD101 était l'étude de la variabilité hydrologique, dynamique, et physico-chimique des grandes masses océaniques de l'océan Atlantique, en liaison avec la climatologie globale et l'évolution des calottes de glace. Le but de cette étude était d'allonger les séries obtenues au cours des campagnes PALEOCINAT et des campagnes équivalentes étrangères, afin de mieux quantifier le rôle du forçage astronomique (et ses composantes autour de 20 et 40 ka) dans l'évolution du climat, tout en prenant en compte l'évolution non linéaire du système interactif formé par les calottes de glace, les mers polaires et la circulation profonde.

Le trajet principal de la campagne suit grossièrement la grande boucle de circulation thermohaline dans l'Atlantique nord: les bordures nord et ouest de l'Atlantique nord permettant d'étudier l'histoire de la NADW (North Atlantic Deep Water), et la bordure est permettant d'aborder les relations avec les eaux profondes antarctiques (Antarctic Bottom Water, AABW) et l'eau intermédiaire de Méditerranée. Des profils verticaux le long des marges et des reliefs sous marins ont été réalisés pour reconstituer la distribution verticale des masses d'eau. Une

grande partie des carottes a été prélevée sur les marges et les rides sédimentaires où les vitesses de sédimentation supérieures à 10 cm/ 1000 ans permettent d'atteindre une résolution temporelle suffisante pour les études des processus climatiques (du siècle à quelques siècles) et permettre ainsi de préciser les relations de phase entre évolution de l'hydrologie de surface et réponse des océans profonds. De tels taux d'accumulations conduisent toutefois généralement à des systèmes sédimentaires instables, et une reconstitution précise de la structure des sédiments avec sondeurs multi-faisceaux et sismique 3.5 khz est nécessaire pour bien contraindre le choix des sites de carottage.

**b-Plusieurs objectifs annexes sont aussi abordés:**

b1- La variabilité haute résolution de l'aimantation rémanente des sédiments au cours des derniers 800 000 ans.

b2 : la variabilité du  $\delta^{18}\text{O}$  des eaux interstitielles et la mémoire de l'évolution du  $\delta^{18}\text{O}$  de l'océan au cours des derniers 150 000 ans.

b3 : l'évolution de la sédimentation dans le bassin nord-ouest de l'Atlantique et la canal profond du NAMOC, en liaison avec l'histoire de la calotte Laurentide et des chenaux fluviaux sous glaciaires.

## **2/ Stratégie d'approche et méthodes:**

La stratégie d'approche a été celle définie par le Groupe WG 100 du SCOR pour IMAGES:

**Lors de la campagne** - Les sites de la campagne IMAGES-MD101 ont été présélectionnés au niveau international en se basant sur l'étude de carottes existantes et/ou des données sismiques 3.5 khz, ou parce que ces sites présentaient un intérêt potentiel exceptionnel pour le programme IMAGES. Lors de la campagne MD101, un court survol de chaque zone présélectionnée a permis de préciser les localisations de stations en analysant la structure des dépôts à l'aide d'un sondeur 3.5 khz. Un repérage au multi-faisceaux des zones sélectionnées devait être également effectué mais le système multi-faisceaux novateur qui est développé par Thompson sur le *Marion Dufresne* n'a pas été opérationnel à temps pour la mission. Plusieurs opérations différentes étaient prévues à chaque site: collecte des sédiments de surface et des eaux de fond, carottage gravité grand volume (1 à 3 m), carottage Kullenberg 20 m doublé par un carottage de 40 à 60 m si les sédiments le permettaient. En fait, pour gagner du temps, l'objectif de carottage de grande longueur a été privilégié. A bord, une batterie d'analyse en continu a été réalisée sur carottes ouvertes ou non, suivant le modèle établi lors des campagnes de l'Ocean Drilling Program (ODP): paramètres physiques (ondes-p, absorption  $\gamma$ , réflectance couleur, susceptibilité magnétique, images digitales), micropaléontologie, sédimentologie. Les équipements d'analyse ont été fournis par les différents laboratoires participants, et leurs résultats comparés en temps réel. L'acquisition par l'IFRTP, en dernière minute, d'une centrale d'acquisition en continu des propriétés physiques (GEOTEK) a permis de réaliser ces études avec le standard de qualité des campagnes ODP. Pendant la campagne IMAGES-MD101, 43 carottes ont été recueillies, parmi lesquelles 29 ont été ouvertes et décrites à bord et 5 ont été échantillonnées en continu (pas d'échantillonnage de 1 cm).

**En laboratoire** - Les paramètres habituels de la paléoclimatologie quantitative seront analysés par les équipes de pointe dans le domaine au niveau international : géochimie et composition isotopique des foraminifères, fonctions de transfert sur les foraminifères et autres micro-fossiles, bio-marqueurs (Uk37), propriétés magnétiques, études des minéraux transportés par les glaces, des cendres volcaniques et des argiles, granulométrie, datations AMS  $^{14}\text{C}$  et, si possible,  $\text{Ar}^{39}/\text{Ar}^{40}$ . Pour les carottes situées à proximité des côtes, les pollens deviendront aussi un paramètre important.

**Règles adoptées pour l'échantillonnage** - Les moitiés archives d'au moins une carotte par station (ou groupe de station) resteront à Gif pour références futures. Chaque investigateur principal (P.I.) co-finançant a accès à un échantillonnage en continu (ex. U-channel) pour chaque projet approuvé par le Comité Scientifique IMAGES. Le volume de chaque échantillonnage en continu dépendra des études programmées (depuis les U-channels 2x2cm jusqu'à une moitié complète de carotte (12.5 cm). Les enregistrements continus non destructifs (imagerie couleur, MST) seront accessibles à tous. Le Comité Scientifique IMAGES est responsable de l'élaboration des coopérations et de la coordination internationale en liaison avec L. Labeyrie, le responsable de projet. F. Bassinot coordonne l'ensemble des échanges entre scientifiques pour l'exploitation de la campagne.

### **3/ Objectifs détaillés:**

**Zone 1 : pente continentale au débouché de la Manche (zones du replat de Trevelyan à 3788 m, MD 95-2001, et du replat de Meriadzec, 2174 m, MD 95-2002).** Sites de test du matériel de carottage, à la sédimentation exceptionnellement rapide (de l'ordre de 50 cm/kyr pendant le stade isotopique 3). Ces sites sont donc très bien placés pour l'étude de la variabilité climatique et sédimentaire lors de la dernière période glaciaire.

**Zone 2 : ride de Feni à l'est du plateau de Rockall (55° N 15° W).** Sur le trajet de la branche initiale de NADW vers le sud et l'ouest, et des eaux de la dérive nord Atlantique quand elles remontent vers la Mer de Norvège. De nombreuses carottes de 5 à 10 m ont été étudiées sur cette ride à la sédimentation rapide mais assez irrégulière. Zone de haute priorité pour CFR-DGO et les co-associés Allemands. De l'ordre de 5 carottes étaient prévues, mais, compte tenu du retard, seulement 2 ont été réalisées (MD 95-2003 et 2004).

**Zone 3: nord ouest de l'Ecosse.** Echantillonnage de zones côtières à très haute vitesse de sédimentation pour l'étude à résolution centennale de l'évolution climatique depuis le dernier maximum glaciaire: Barra Fan, : Lat. 57°018N, 10°035 W, 2000 m (MD 95-2005 - en dehors du site optimal- et MD 95-2006) et St Kilda Basin: 57°31'N, 08°23'W 158m (MD 95-2007). Objectifs de l'Université d'Edinburg (W. Austin).

**Zone 4 : Ride des îles Féroé.** Zone déjà étudiée par des carottes courtes. Intérêt exceptionnel, car on retrouve enregistrées à la fois les oscillations de la pénétration de l'eau de surface nord Atlantique, et la réponse des eaux profondes au seuil vers l'Atlantique nord. 62°58N 2°22W 1760m. Vitesse de sédimentation environ 10 cm/ka. Zone de haute priorité pour le groupe CFR-DGO et les équipes Allemandes. 2 carottes réalisées (MD 95-2008 et MD 95-2009).

**Zone 5 : Plateau de Vöring.** Zone déjà connue pour l'étude de la période récente par des carottages courts, et pour la partie ancienne par le forage ODP 644. Intérêt exceptionnel pour suivre l'évolution de la calotte de glace scandinave en liaison avec les enregistrements du Groenland (GRIP et GISP) et l'évolution de la circulation thermohaline en Mer de Norvège. Vitesse de sédimentation rapide. Possibilité d'étendre la couverture existante en fonction de la proximité de la Norvège et de la profondeur d'eau. Sites réalisés MD 95-2010 66°58N, 07°38 E, 1048 m ; MD 95-2011 66°41N, 04°34 E, 1226 m. Priorité pour l'Univ. de Bergen (E. Jansen) et les équipes Allemandes.

**Zone 6 : Bear Island.** Cette zone, située en avant de la petite calotte de la Mer de Barentz lors de la dernière période glaciaire, joue un rôle exceptionnel dans la compréhension des processus liés au début des déglaciations. Le carottage d'une zone d'accumulation à forte vitesse de sédimentation constituait l'objectif prioritaire de l'équipe de l'Univ. de Kiel. Les carottes ont été malheureusement assez perturbées lors du prélèvement (MD 95-2012 72°09N, 11°26E, 2094m et MD 95-2013 72°11N, 13°06E, 1638 m).

**Zone 7 : marge sud de l'Islande (Reykjanes, 60°N-22°W).** C'est une zone explorée lors de la campagne Paleocinat 1 du N/O Suroît (carottes SU 90-32 et 33, 10 à 15 m de long à 8-12 cm/ka), très favorable à l'étude sur plusieurs cycles glaciaires-interglaciaires de la remontée vers le nord et la Mer de Norvège des eaux chaudes de la dérive nord Atlantique, les liens avec l'évolution de la calotte Islandaise, et l'évolution des eaux intermédiaires à partir de la composition isotopique des foraminifères benthiques. Nombreux niveaux de cendres volcaniques datables en Ar<sup>39</sup>-Ar<sup>40</sup>, quand la méthode sera fonctionnelle au CFR, pour améliorer la chronologie absolue dérivée de la stratigraphie isotopique. Objectif prioritaire CFR-DGO. Une carotte réalisée à l'emplacement de la carotte SU 90-33 (MD 95-2014 60°35 N, 22°05W, 2397 m).

**Zone 8 : Ride de Gardar et ride médio-Atlantique (55°-57°N-30-40°W).** Les derniers travaux du CFR et du DGO semblent montrer que c'est dans cette zone que se forment les eaux profondes en période glaciaire, par un mécanisme voisin de celui se déroulant actuellement en Mer du Labrador (eaux modales). Il n'existe pas de bonne carotte étudiée sur la zone. Objectif prioritaire CFR-DGO. Deux carottes ont été réalisées, MD 95-2015 à 58°46 N 25°58 W 2630 m et MD 95-2016 à 57°43 N et 29°26 W 2318 m. La seconde de ces carottes a été prélevée sur un site déjà échantillonné par W. Ruddiman, pour lequel la vitesse de sédimentation en période interglaciaire est exceptionnelle (supérieure à 30 cm/kyr).

**Zones 9-10 : Marge nord est du continent Américain (44°-50°N : 40°-55°W).** C'est la zone qui doit enregistrer de façon la plus sensible les arrivées d'icebergs associées aux débâcles des "niveaux de Heinrich" issus de la région nord de la Laurentide. En profondeur, on peut y enregistrer les caractéristiques des eaux profondes (NADW) dans leur trajet vers le sud, hors des zones de formation. Plusieurs zones carottées par le Lamont (LDEO, Univ. de Columbia, NY), les canadiens ou lors de la campagne Paléocinat 1, ont servi de cible. Cette zone est l'objectif unique des équipes canadiennes. Un autre objectif, lui aussi Canadien, vise un

échantillonnage serré d'une portion du canyon du NAMOC. Ce canyon suit pendant presque 2000 km la pente entre le talus continental du Labrador et les profondeurs du bassin ouest Atlantique. Le canyon est apparemment creusé par des masses d'eau turbides issues des écoulements sous glaciaires lors des périodes d'extension de la calotte Laurentide. 8 carottes, dont l'une de 34.8 m, ont été réalisées sur cette zone (MD 95-2018 à 23). La première des carottes pour le programme IMAGES Canadien a été réalisée en bordure de la fracture de Gibbs, sur la ride médio-Atlantique au sud-est de Gardar (MD 95-2017 53° 03 N, 33°32W, 3100 m). Les autres carottes ont été réalisées sur la marge Canadienne au nord, à l'est et au sud-ouest de Terre Neuve (MD 95 2024 à 2033), entre 4150 m et 878 m de profondeur.

Zone 11 : Ride des Bermudes. Zone de grand intérêt pour suivre à très haute résolution (vitesse de sédimentation de 30 cm/ka ou plus) les oscillations, en surface, des eaux subtropicales du Gulf Stream, et en profondeur la limite entre NADW et AABW. Trois carottages successifs ont été réalisés afin de couvrir les derniers cycles climatiques (MD95-2034 à MD95-2036). La carotte MD95-2036 fait 52.64 m de long, le record de longueur pour un carottage Kullenberg pour l'Atlantique). Principal objectif des équipes US.

Zones 12-13: Ride médio-Atlantique, Açores et Marge ouest de la Péninsule Ibérique (37°N 10°W). La zone comprenant la marge ibérique et la zone des Açores, étudiée en particulier par les campagnes de 1980 et 1990 du Suroît et des campagnes du Météor, présente une forte vitesse de sédimentation (entre 10 et 50 cm/1000 ans). Elle permet de suivre l'évolution de la bordure sud du front polaire en période glaciaire, et permet de reconstituer, à partir des foraminifères benthiques, la contribution des eaux de la Méditerranée aux eaux intermédiaires Atlantiques. Les carottes du pourtour des Açores offrent des enregistrements paléovolcaniques et paléomagnétiques d'un intérêt exceptionnel. Ces zones sont prioritaires pour les équipes du LGQ, de l'URA Orsay-Muséum, de l'IFREMER à Brest, du LODYC-Paris, du groupe CFR-DGO, les équipes allemandes (GEOMAR) et portugaises (Portugal Geol. Surv.), et les équipes anglaises de l'IOS et de l'Université de Cambridge. De 8 à 10 carottes de très grandes longueurs étaient prévues sur ces zones, mais seulement 6 ont pu être réalisées, faute de temps disponible (MD 95-2037 - 2042).

Zone 14 : Bordure ouest de la Méditerranée: Zone de forte accumulation où il sera possible de suivre l'évolution paléoclimatique de la bordure sud de l'Europe (par pollens et bio-marqueurs) en liaison avec l'hydrologie méditerranéenne. Priorité de l'équipe Univ. de Barcelone et de Salamanque (et collaboration avec Brest -Ifremer) : MD 95-2043 à 36°08 N, 02°37W, 1841 m).



Date	Station	GMT Time	Latitude (N)	Longitude (W)	Water depth (m)	Operation
05/29/95		23h00				Departure from Brest for station I
05/30/95	I	11h20 14h59 19h30	46°48.17	08°40.28	3788	arrived on station (Trevelyan) Calypso (22.37 m; MD 95-2001) departure for station II
	II	20h55 23h07	47°27.12	08°32.03	2174	arrived on station (Mériadzec) Calypso (29.99 m; MD 95-2002)
05/31/95		00h50				departure for station III
06/01/95	III	11h00 12h07 14h52	55°10.61	14°45.32	2365	arrived on station (Feni Drift) Calypso (26.93 m; MD 95-2003) departure for station IV
	IV	19h30 20h55 22h54	55°28.07	14°40.51	2177	arrived on station (Feni Drift) Calypso (28.485 m; MD 95-2004) departure for station V
06/02/95	V	10h02 11h15	57°02	10°03.83	2130	arrived on station (Barra Fan) Calypso (22.02 m; MD 95-2005) boat drifting, core taken 2 miles from the proposed site
		12h36 17h17 17h55	57°01.81	10°03.48	2122	return to proposed site Calypso (30.22 m; MD 95-2006) departure for station VI
06/03/95	VI	23h20				arrived on station (St-Kildas basin)
06/04/95		00h03 00h51	57°31.06	08°23.17	158	Calypso (19.35 m; MD95-2007) departure for station VII
	VII	13h35 14h08	62°44.35	03°59.64	1016	arrived on station (Faeroes) Calypso (21.5 m; MD 95-2008)
06/05/95		06h51 10h30	62°44.25	03°59.86	1027	Dynamic Positioning System test Calypso (32.43 m; MD 95-2009) departure for Faeroes port call, station VIII
06/06/95	VIII	09h48 16h00	61°59.75	06°45.03	<100	arrived at the Faeroes port call, station VIII Gravity core (MD 9508GR01)
06/07/95	IX	20h30 21h27 22h50	66°41.05	04°33.97	1226 100	arrived on station (Vöring plateau) Calypso (32.53 m; MD 95-2010) plankton tow (MD 95-IX-2)
06/08/95		01h10				departure for station X
06/08/95	X	06h45 07h30 09h06	66°58.18	07°38.36E	1048	arrived on station (Vöring Plateau) Calypso (17.49 m; MD 95-2011) departure for station XI
06/09/95	XI	05h42 07h11 08h24 09h00	72°09.06	11°26.06E	2094 100	arrived on station (Bear Island Fan) Calypso (19.76 m; MD 95-2012) plankton tow (MD 95-XI-3) departure for station XII

Date	Station	GMT Time	Latitude (N)	Longitude (W)	Water depth (m)	Operation
	XII	12h15 13h06 14h45	72°10.92	13°05.63E	1638	arrived on station (Bear Island slope) Calypso (13.10 m; MD 95-2013) departure for Iceland port call- station XIII
06/11/95	XIII	19h22 19h25 21h50	64°05.40	15°15.73	100	arrived at the Iceland port call (Hornfjordur) station XIII; Box core (empty) departure for station XIV
06/12/95	XIV	16h10 17h23 19h08 19h32	60°34.93	22°04.52	2397 100	arrived on station (Reykynes Ride) Calypso (33.61 m; MD 95-2014) plankton tow (MD 95-XIV-4) departure for station XV
06/13/95	XV	06h36 08h34 08h00 10h31	58°45.74	25°57.54	2630 100	arrived on station ( Reykynes Ride) Calypso (34.42 m; MD 95-2015) plankton tow (MD-XV-5) departure for station XVI
	XVI	19h50 20h45 21h12	57°42.46	29°25.44	2318	arrived on station (Reykynes) Calypso (32.83 m; MD 95-2016) departure for station XVII
06/14/95	XVII	19h03 20h31	53°02.54	33°31.64	3100	arrived on station (Gibbs Fracture Zone) Calypso; corer lost during removal from the sediment
06/15/95		20h56 02h50 04h45 05h55	53°02.56	33°31.51	100 3100	plankton tow (MD 95-XVII-6) Calypso (34.59 m; MD 95-2017) hydro cast departure for station XVIII
06/16/95	XVIII	08h00 09h29 12h10	50°43.88	43°08.54	4221	arrived on station (Namoc) Calypso (14.70 m; MD 95-2018) departure for station XIX
	XIX	15h03 16h11 17h30	51°09.22	43°17.08	4360	arrived on station (Namoc) Gravity core (empty; smear slide MD 9519GR02) departure for station XX
	XX	18h28 20h03 21h41	51°05.68	43°13.27	4262	arrived on station (Namoc) Calypso (12.86 m; MD 95-2019) departure for station XXI
06/17/95	XXI	23h00 00h30 01h51	51°02.42	43°00.62	4289	arrived on station (Namoc) Calypso (8.63 m; MD 95-2020) departure for station XXII
	XXII	03h40 05h15 06h45	50°59.76	42°55.19		arrived on station (Namoc) Gravity core (1.21 m; MD 9522GR03) "petit-chef" first gravity core !!! departure for station XXIII

Date	Station	GMT Time	Latitude (N)	Longitude (W)	Water depth (m)	Operation
06/18/95	XXIII	07h30				arrived on station (Namoc)
		09h00	50°51.81	42°44.49	4283	Calypso (4.48 m; MD 95-2021)
		10h50				departure for station XXIV
	XXIV	14h24				arrived on station (Namoc)
		15h43	50°34.38	43°03.51	4245	Calypso (18.37 m; MD 95-2022)
		16h14 17h23			100	plankton tow (MD 95-XXIV-7) departure for station XXV
	XXV	20h41				arrived on station (Namoc)
		23h48	50°58.32	43°13.04	4198	Calypso (34.68 m; MD 95-2023)
		01h50				departure for station XXVI
	XXVI	10h00				arrived on station (Orphan Knoll)
		11h11	50°12.40	45°41.22	3539	Calypso (29.52 m; MD 95-2024)
		13h24				departure for station XXVII
XXVIIa	13h45				arrived on station (Orphan Basin)	
	18h31				core touched the bottom after releasing in the water	
	18h34	49°47.41	46°49.03		core-catcher recovered in one bag (MD 95-XXVII-8; Attention: samples MD 95-XXVII-8 was renumbered after having registered MD 95-XXVII-7)	
06/19/95	XXVIIb	00h20 02h13	49°47.65	46°41.85	3009	Calypso (35.12 m; MD 95-2025) departure for station XXVIII
XXVIII	05h45				arrived on station (Orphan Basin)	
	09h16	50°12.03	45°41.33	3539	Calypso (bent; empty) recovered core-catcher in one bag (MD 95-XXVIII-9; Attention: samples MD 95-XXVIII-9 was renumbered after having registered MD 95-XXVIII-7)	
	12h45				departure for station XXIX	
06/20/95	XXIX	23h11				arrived on station (Sackville ridge)
		23h47	48°13.62	46°39.39	878	Calypso (27.91 m; MD 95-2026)
		23h40 01h10			100	plankton tow (MD 95-XXIX-10) departure for station XXX
06/21/95	XXXa	05h49				arrived on station (S-E of Newfoundland)
		08h07 (?)	41°45.58	47°19.94		no visible release on the tensionmeter recording stones in the core-catcher, recovered in one bag (MD 95-XXX-11)
XXXb	15h30	41°44.67	47°24.79	4112	Calypso (47.67 m; MD 95-2027)	
	17h20				departure for station XXXI	
06/22/95	XXXI	10h22				arrived on station (Fogo Seamount)
		11h37	42°05.99	52°44.85	3368	Calypso (34.20 m; MD 95-2028)
		13h20				departure for station XXXII
XXXII	22h20				arrived on station (Laurentien Fan)	
	22h49	43°06.78	55°15.96	4156	Calypso (35.10 m; MD 95-2029)	

Date	Station	GMT Time	Latitude (N)	Longitude (W)	Water depth (m)	Operation
06/23/95		01h55				departure for station XXXIII
	XXXIII	11h00				arrived on station (Narwhal Area)
		12h41	44°19.14	53°43.44	1573	Calypso (35.59 m; MD 95-2030)
						core in very bad condition, redo
	XXXIV	21h47	44°18.45	53°44.14	1570	Calypso (27.72 m; MD 95-2031)
		23h00				departure for station XXXV
06/24/95	XXXV	05h32				arrived on station ( Laurentien margin)
		07h06	44°39.46	55°37.14	1433	Calypso (28.54 m; MD 95-2032)
						departure for St-Pierre et Miquelon
		19h00				Arrived at the St-Pierre et Miquelon port call
06/26/95		08h24				Departure from St-Pierre et Miquelon for station XXXVI
	XXXVI	19h50				arrived on station (Laurentien margin)
		20h53	44°39.87	55°37.21	1412	Calypso (29.68 m; MD 95-2033)
		23h36			100	plankton tow (MD 95-XXXVI-12)
06/27/95		00h00				departure for station XXXVII
06/28/95	XXXVII	19h53				arrived on station (Bermuda Rise)
		21h24	33°41.44	57°34.55	4461.5	Calypso (47.16 m; MD 95-2034)
		21h53			100	plankton tow (MD 95-XXXVII-13)
06/29/95		00h05				departure for station XXXVIII
	XXXVII	04h01				arrived on station (Bermuda Rise)
		05h34	33°29.20	57°53.16	4286	Calypso (39.25 m; MD 95-2035)
		09h00				departure for station XXXIX
	XXXIX	12h14				arrived on station (Bermuda Rise)
		14h43	33°41.44	57°34.55	4461.5	Calypso (52.64 m; MD 95-2036)
		16h47				departure for station XL
07/03/95	XLa	02h38				arrived on station (South Lucky)
			37°05.22	32°01.82	~1800	Calypso; Kevlar cable broke; corer, piston and cable (~1800 m) lost on the bottom
	XLb	11h16	37°05.23	32°01.87	2159	Calypso (35.81 m; MD 95-2037)
		11h52			100	plankton tow (MD 95-XL-14)
		14h05				departure for station XLI
07/04/95	XLI	07h25				arrived on station
		08h26	37°45.15	27°11.21	2310	Calypso (37.85 m; MD 95-2038)
		10h25				departure for the Azores port call
		16h00				arrived at the Azores port call
		22h40				departure from Azores for station XLII
07/06/95	XLII	23h15				arrived on station
07/07/95		00h17	40°34.71	10°20.91	3381	Calypso (35.71 m; MD 95-2039)

Date	Station	GMT Time	Latitude (N)	Longitude (W)	Water depth (m)	Operation
		02h00				departure for station XLIII
07/07/95	XLIII	05h34				arrived on station
		06h52	40°34.91	09°51.67	2465	Calypso (35.24 m; MD 95-2040)
		08h15				departure for station XLIV
07/08/95	XLIVa	20h05				arrived on station
		21h18	37°49.97	09°30.62	1123	Calypso (piston not released)
	XLIVb	22h58	37°50	09°30.65	1123	Calypso (17.62 m; MD 95-2041)
07/09/95		00h00				departure for station XLV
		04h45				arrived on station
			37°48	10°09.99	3146	Calypso (39.56 m; MD 95-2042)
		08h10				departure for station XLVI
		09h10				arrived on station
		09h59	36°08.60	02°37.27	1841	Calypso (37.46 m; MD 95-2043)
		11h00				departure for Marseille



Date	Station	Water depth (m)	Latitude (N)	Longitude (W)	Location	Core #	Length (m)	Sections	Comments
05/30/95	I	3788	46°48.17	08°40.28	Trevelyan	MD 95-2001	22.37	XVI	<b>Giant Kullenberg core (Calypso)</b> **liner exploded between 0-533 cm; 3.5 Khz record was partially erased
05/30/95	II	2174	47°27.12	08°32.03	Meriadzec	MD 95-2002	29.99	XX	<b>Calypso core</b> *** OK
06/01/95	III	2365	55°10.61	14°45.32	Feni Drift	MD 95-2003	26.93	XVIII	<b>Calypso core</b> **Liner exploded on the upper 8m approx.; very perturbed sediments in sections I to IV (0-1050 cm) and section XIX (2683-2715 cm)
06/01/95	IV	2177	55°28.07	14°40.51	Feni Drift	MD 95-2004	28.485	XIX	<b>Calypso core</b> *** OK
06/02/95	V	2130	57°02	10°03.83	Barra fan	MD 95-2005	22.02	XV	<b>Calypso core</b> **boat drifting, core taken 2 miles from the proposed site; redo (MD 95-2006); numerous voids in the core
		2122	57°01.81	10°03.48	Barra fan	MD 95-2006	30.22	XX	<b>Calypso core</b> ***
06/04/95	VI	158	57°31.06	08°23.17	St Kildas Basin	MD 95-2007	19.35	XIII	<b>Calypso core</b> *** stopped on stones
06/04/95	VII	1016	62°44.35	03°59.64	Nord Faeroes	MD 95-2008	21.5	XVII	<b>Calypso core</b> ** liner sliced along from sections XIII to XV (1715-2075 cm); holocene missing from the top
		1027	62°44.25	03°59.86	Nord Faeroes	MD 95-2009	32.43	XXI	<b>Calypso core</b> **upper part of the core bent; liner sliced along between about 1330-2224 cm
06/06/95	VIII	< 100	61°59.75	06°45.03		MD 9508GR01		bag	<b>Gravity core</b> about 1 kg of mud and shells
06/07/95	IX	1226	66°41.05	04°33.97	Vöring plateau	MD 95-2010	32.53	XXI	<b>Calypso core</b> ** OK
		100				MD 95-IX-2			<b>plankton tow</b> (Attention: no sample n°1 -> gravity core from the site VIII was renumbered MD9508GR01 after having registered as a sample (MD 95-VIII-1))
06/08/95	X	1048	66°58.18	07°38.36E	Vöring plateau	MD 95-2011	17.49	XII	<b>Calypso core</b> Site location wrong, slumps. Bent core; liner imploded in the middle of the core; 30 cm of sediment of the top recovered in a 1/2 liner

Date	Station	Water depth (m)	Latitude (N)	Longitude (W)	Location	Core #	Length (m)	Sections	Comments
06/09/95	XI	2094	72°09.06	11°26.06E	Bear Isl. Fan	MD 95-2012	19.76	XV	<b>Calypso core</b> * trigger release jam, first touch without releasing; sent second messenger -> double touch. Bent core; broken liner; liner overlapped in section VI; between 1702-1976 cm, sediments extruded in the core and recovered in a liner plankton tow
		100				MD 95-XI-3			
06/09/95	XII	1638	72°10.92	13°05.63E	Bear Isl. Fan	MD 95-2013	13.1	IX	<b>Calypso core</b> bent core; sliced liner all along the core
06/11/95	XIII	100	64°05.40	15°15.73			0	0	<b>Gravity core</b> empty
06/12/95	XIV	2397	60°34.93	22°04.52	S. Iceland	MD 95-2014	33.61	XXIII	<b>Calypso core</b> *** OK; about 7 cm of sediment dropped on the deck when cutting between sections plankton tow
		100				MD 95-XIV-4			
06/13/199	XV	2630	58°45.74	25°57.54	Reykjanes Ridge	MD 95-2015	34.42	XXIII	<b>Calypso core</b> *** OK
		100				MD 95-XV-5			<b>plankton tow</b>
06/13/199	XVI	2318	57°42.46	29°25.44	Reykjanes Ridge	MD 95-2016	32.83	XXII	<b>Calypso core</b> *** OK; broken liner in sections I, II and VII; lost of liquid sediment at the top.
06/14/199	XVII	3100	53°02.54	33°31.64	Gibbs				<b>Calypso core</b> core lost during removal from the sediment, piston cable broke at 6 tons plankton tow
		100				MD 95-XVII-6			
06/15/199		3100	53°02.56	33°31.51	Gibbs fracture	MD 95-2017	34.59	XXIV	<b>Calypso core</b> * supply expanded sediment in sections I to VIII (27-1200 cm); section XVI perturbed with many liquid pockets hydro cast
		133							
06/16/199	XVIII	4221	50°43.88	43°08.54	Namoc	MD 95-2018	14.7	X	<b>Calypso core</b> Sections I to IVa (0-525 cm) damaged and mostly empty; 140 cm of sediment missing between sections VI and VIII

Date	Station	Water depth (m)	Latitude (N)	Longitude (W)	Location	Core #	Length (m)	Sections	Comments
06/16/199	XIX	4360	51°09.22	43°17.08	Namoc	MD 9519GR02	0	0	Gravity core empty; smear slide
06/16/199	XX	4262	51°05.68	43°13.27	Namoc	MD 95-2019	12.86	IX	Calypso core *** OK
06/17/1995	XXI	4289	51°02.42	43°00.62	Namoc	MD 95-2020	8.63	VI	Calypso core * upper 1-2m imploded; 20 cm of the sediment from the top was put in a liner (1a), the rest of the sediment are in one bag (1b), regular core cutting starts with section 1c (0-150 cm)
06/17/1995	XXII	4415	50°59.76	42°55.19	Namoc	MD 9522GR03	1.21	1	Gravity core ** first touch at very low speed in order to spot the bottom, core entered at 2 m/s;
06/17/1995	XXIII	4283	50°51.81	42°44.49	Namoc	MD 95-2021	4.48	III	Calypso core core cutter lost but core-catcher recovered, gravels at the base; upper part of the liner imploded above the sediment; top 0-30 cm recovered in one bag (1a); 30-150 cm interval is half full (section 1b)
06/17/1995	XXIV	4245 100	50°34.38	43°03.51	Namoc	MD 95-2022 MD 95-XXIV-7	18.37	XIII	Calypso core * liner imploded at mid-pipe (~6m) ; overlapping of 3 liners in section IV; liquid plankton tow
06/17/1995	XXV	4198	50°58.32	43°13.04	Namoc	MD 95-2023	34.68	XXIV	Calypso core ** liquid pocket between sections IXb-X, partially recovered in one bag
06/18/1995	XXVI	3539	50°12.40	45°41.22	Orphan Knoll	MD 95-2024	29.52	XX	Calypso core ** liquid pocket between sections X-XI, partially recovered in one bag
06/18/1995	XXVIIa	2916	49°47.41	46°49.031	Orphan Basin	MD 95-XXVII-8	0	0	Calypso core the core touched the bottom after releasing in the water; core-catcher recovered in one bag (stones in core-catcher 1-3 cm diameter); Attention: samples MD 95-XXVII-8 was renumbered after having been registered MD 95-XXVII-7
06/19/199	XXVIIb	3009	49°47.646	46°41.85	Orphan Basin	MD 95-2025	35.12	XXIV	Calypso core *** OK; liquid pocket between sections IX-X

Date	Station	Water depth (m)	Latitude (N)	Longitude (W)	Location	Core #	Length (m)	Sections	Comments
06/19/199	XXVIII	3539	50°12.03	45°41.33	Orphan Knoll	MD 95-XXVIII-9	0	0	<b>Calypso core</b> bent and empty core; stones above the core-catcher; Attention: samples MD 95-XXVIII-9 was renumbered after having registered MD 95-XXVIII-7
06/19/199	XXIX	878	48°13.62	46°39.39	Sackville spur	MD 95-2026	27.91	XIX	<b>Calypso core</b> bent core; liquid pocket between sections V-VI; core catcher full of stones
		100				MD 95-XXIX-10			<b>plankton tow</b>
06/21/199	XXXa	4095	41°45.58	47°19.94	Newfoundland Slope	MD 95-XXX-11	0	0	<b>Calypso core</b> no visible release on the tensionmeter recording; empty core; stones in the core-catcher, recovered in one bag
06/21/199	XXXb	4112	41°44.67	47°24.79	Newfoundland Slope	MD 95-2027	47.67	XXXII	<b>Calypso core</b> *** OK
06/22/199	XXXI	3368	42°05.99	52°44.85	Fogo Sea Mount	MD 95-2028	34.2	XXIII	<b>Calypso core</b> *** OK; liquid pocket between sections VII-VIII (very fluid), partially recovered in
06/22/199	XXXII	4156	43°06.78	55°15.96	Laurentian fan	MD 95-2029	35.1	XXIV	<b>Calypso core</b> ** OK; liquid pockets partially recovered between sections IX-X and XI-XII; section I partially emptied during handling on deck
06/23/199	XXXIII	1573	44°19.14	53°43.44	Narwhal	MD 95-2030	35.59		<b>Calypso core</b> core in very bad condition, redo
06/23/199	XXXIV	1570	44°18.45	53°44.14	Narwhal	MD 95-2031	27.72	XIX	<b>Calypso core</b> ** OK; core slightly bent at the top (~5m); broken liner in sections VI and VII
06/24/199	XXXV	1433	44°39.46	55°37.14	upper Laurentian sl.	MD 95-2032	28.54	XX	<b>Calypso core</b> liner bent at the top (0-600 cm (sections I to IV)); upper 3 sections ± sliced
06/26/199	XXXVI	1412	44°39.87	55°37.21	upper Laurentian sl.	MD 95-2033	29.68	XX	<b>Calypso core</b> ** OK; ore slightly bent at the top; when extruding the liner, 70 cm of sediment fell on deck, recovered in a 1/2 liner (714-784 cm)
		100	44°39.15	55°36.50		MD 95-XXXVI-			
06/28/199	XXXVII	4461.5	33°41.46	57°34.55	Bermuda Rise	MD 95-2034	47.16	XXXII	<b>Calypso core</b>

Date	Station	Water depth (m)	Latitude (N)	Longitude (W)	Location	Core #	Length (m)	Sections	Comments
		100	33°41.54	57°35.30	Bermuda Rise	MD 95-XXXVII-			*** OK; sections I to III very fluid; liner imploded in sections IX and X; bags of fluids sediments recovered between sections XIII-XIV and XVI-XVII <b>plankton tow</b>
06/29/199	XXXVII	4286	33°29.20	57°53.16	Bermuda Rise	MD 95-2035	39.25	XXVII	<b>Calypso core</b> *** OK; first touch without releasing, released on the second attempt
06/29/199	XXXIX	4461.5	33°41.44	57°34.55	Bermuda Rise	MD 95-2036	52.64	XXXVI	<b>Calypso core</b> *** OK; first touch without releasing, released on the second attempt; liner imploded in sections II and III; sections III et IV very fluid; one bag of fluid sediment recovered between sections XIV and XV
03/07/95	XLa	1800	37°05.22	32°01.82	AZORES/Sud Lucky		0	0	<b>Calypso core</b> piston cable broke; piston, corer and 1800 m of cable lost....
03/07/95	XLb	2159	37°05.23	32°01.87	AZORES/Sud Lucky	MD 95-2037	35.81	XIII	<b>Calypso core</b> *** OK; upper part of the core bent; 0-4 cm disturbed during core cutting; lost of sediment between 283-336 cm (section IIb) <b>plankton tow</b>
		100				MD 95-XL-14			
07/04/95	XL1	2310	37°45.15	27°11.21	AZORES	MD 95-2038	37.85	XXVI	<b>Calypso core</b> *** OK; many turbidites
07/07/95	XLII	3381	40°34.71	10°20.91	Porto Seamount	MD 95-2039	35.71	XXIV	<b>Calypso core</b>
07/07/95	XLIII	2465	40°34.91	09°51.67	Porto Seamount	MD 95-2040	35.24	XXIV	<b>Calypso core</b> core bent; section IIb jamed in the bent part of the core, recuperated in 1/2 liner
07/08/95	XLIVa	1123	37°49.97	09°30.62					<b>Calypso core</b> piston not released
07/08/95	XLIVb	1123	37°50	09°30.65	marge ibérique	MD 95-2041	17.62	XII	<b>Calypso core</b> upper part of the core bent; liner exploded in section Ia
07/09/95	XLV	3146	37°47.99	10°09.99	marge ibérique	MD 95-2042	39.56	XXVII	<b>Calypso core</b> ** OK; core bent
07/09/95	XLVI	1841	36°08.60	02°37.27	Alboran Sea	MD 95-2043	37.46	XXV	<b>Calypso core</b> *** OK; liner exploded in section VIII

Cores #	Water depth (m)	Latitude (N)	Longitude (W)	Length (m)
MD 95-2001	3788	46°48.17	08°40.28	22.37
MD 95-2002	2174	47°27.12	08°32.03	29.99
MD 95-2003	2365	55°10.61	14°45.32	26.93
MD 95-2004	2177	55°28.07	14°40.51	28.485
MD 95-2005	2130	57°02	10°03.83	22.02
MD 95-2006	2122	57°01.81	10°03.48	30.22
MD95-2007	158	57°31.06	08°23.17	19.35
MD 95-2008	1016	62°44.35	03°59.64	21.5
MD 95-2009	1027	62°44.25	03°59.86	32.43
MD 95-2010	1226	66°41.05	04°33.97	32.53
MD 95-2011	1048	66°58.18	07°38.36E	17.49
MD 95-2012	2094	72°09.06	11°26.06E	19.76
MD 95-2013	1638	72°10.917	13°05.632E	13.1
MD 95-2014	2397	60°34.93	22°04.52	33.61
MD 95-2015	2630	58°45.74	25°57.54	34.42
MD 95-2016	2318	57°42.46	29°25.44	32.83
MD 95-2017	3100	53°02.56	33°31.51	34.59
MD 95-2018	4221	50°43.88	43°08.54	14.7
MD 95-2019	4262	51°05.68	3°13.27	12.86
MD 95-2020	4289	51°02.42	43°00.62	8.63
MD 95-2021	4283	50°51.81	42°44.49	4.48
MD 95-2022	4245	50°34.38	43°03.51	18.37
MD 95-2023	4198	50°58.32	43°13.04	34.68
MD 95-2024	3539	50°12.40	45°41.22	29.52
MD 95-2025	3009	49°47.65	46°41.85	35.12
MD 95-2026	878	48°13.62	46°39.39	27.91
MD 95-2027	4112	41°44.67	47°24.79	47.67
MD 95-2028	3368	42°05.99	52°44.85	34.2
MD 95-2029	4156	43°06.78	55°15.96	35.1
MD 95-2030	1573	44°19.14	53°43.44	35.59
MD 95-2031	1570	44°18.45	53°44.14	27.72
MD 95-2032	1433	44°39.46	55°37.14	28.54
MD 95-2033	1412	44°39.87	55°37.21	29.68
MD 95-2034	4461.5	33°41.46	57°34.55	47.16
MD 95-2035	4286	33°29.20	57°53.16	39.25
MD 95-2036	4461.5	33°41.44	57°34.55	52.64
MD 95-2037	2159	37°05.23	32°01.87	35.81
MD 95-2038	2310	37°45.15	27°11.21	37.85
MD 95-2039	3381	40°34.71	10°20.91	35.71
MD 95-2040	2465	40°34.91	09°51.67	35.24
MD 95-2041	1123	37°50	09°30.64	17.62
MD 95-2042	3146	37°47.99	10°09.99	39.56
MD 95-2043	1841	36°08.60	02°37.27	37.46

Cores #	Water depth (m)	Latitude (N)	Longitude (W)	Length (m)
MD 9508GR01	< 100	61°59.75	06°45.03	0.2
MD 9519GR02	4360	51°09.22	43°17.08	0 (smear slide)
MD 9522GR03	4415	50°59.76	42°55.19	1.21

Samples #	Water depth (m)	Latitude (N)	Longitude (W)	Type of sampling
MD 95-IX-2	100	66°41.05	04°33.97	plankton tow
MD 95-XI-3	100	72°09.06	11°26.06E	plankton tow
MD 95-XIV-4	100	60°34.93	22°04.52	plankton tow
MD 95-XV-5	100	58°45.74	25°57.54	plankton tow
MD 95-XVII-6	100	53°02.54	33°31.64	plankton tow
MD 95-XXIV-7	100	50°34.38	43°03.51	plankton tow
MD 95-XXVII-8*	2916	49°47.41	46°49.03	core-catcher
MD 95-XXVIII-9**	3539	50°12.03	45°41.33	core-catcher
MD 95-XXIX-10	100	48°13.615	46°39.394	plankton tow
MD 95-XXX-11	4095	41°45.58	47°19.94	core-catcher
MD 95-XXXVI-12	100	44°39.15	55°36.50	plankton tow
MD 95-XXXVII-13	100	33°41.54	57°35.30	plankton tow
MD 95-XL-14	100	37°05.23	32°01.87	plankton tow

\* samples MD 95-XXVII-8 was renumbered after having registered MD 95-XXVII-7

\*\* samples MD 95-XXVIII-9 was renumbered after having registered MD 95-XXVIII-7



# ON BOARD MEASUREMENTS

## A. Preliminary report on physical properties measurements

Gavin Dunbar, Franck Bassinot, Kate Jarrett, and the physical properties shipboard scientists

*Several types of physical properties measurements were performed on board the Marion Dufresne during the IMAGES-MD101 cruise. Near-continuous, non destructive measurements of bulk density, compressional wave velocity, and magnetic susceptibility were obtained with a GEOTEK MultiSensor Track (MST). Index properties measurements (bulk density, porosity, water content) were also performed on discrete samples obtained with a 10 cc syringe. The Minolta CM-2002 hand-held spectrophotometer was used to measure reflected visible light in thirty one 10-nm wide bands ranging from 400-700 nm. Measurements were usually performed every 5 cm. Continuous video scanning of the working half sections was performed using the system developed by the CFR.*

*All the physical properties data obtained on board the R/V Marion Dufresne during the MD101 cruise are available on a FTP server at the Centre des Faibles Radioactivités (access to the data base requires a specific user identification and password that are available to people involved in post-cruise studies. Contact: Franck.Bassinot@cfr.cnrs-gif.fr).*

### 1. Continuous measurements of physical properties (MST).


High-resolution logging of physical properties was carried out on rounded cores on board the R/V *Marion Dufresne* using a GEOTEK Ltd Multi Sensor Core Logger (or MST, Multi Sensor Track). These measurements formed the basis for initial correlations between sites. When combined with the shipboard spectrophotometer data and the smear slide descriptions, these data enabled a tentative stratigraphy to be assigned to the cores within hours of recovery. Combined plots of selected MST and spectrophotometer data are shown elsewhere in this report.

#### **1.1 Description of the Multi Sensor Core Logger**


The shipboard multi sensor core logger system have three main sensor units (the configuration of the shipboard system is shown in figure 1):

- 1) a P-WAVE LOGGER that consists of two transducers (piezzo-electric crystals) connected to an oscilloscope to measure the travel time of a 500 kHz compressional wave through the core. The travel time is then used to calculate P-wave velocity in the sediment.
- 2) a GAMMA RAY SOURCE (137Cs) AND DETECTOR for measuring the bulk density by comparing the attenuation of gamma rays through the cores with attenuation through aluminum standards.
- 3) a MAGNETIC SUSCEPTIBILITY METER for determining the amount of magnetically susceptible material present in the cores. In our case, a Bartington Instruments MS2 loop sensor with an internal diameter of 125 mm was used.

In order to calibrate the values derived from these sensors, core temperature and core diameter were measured on each section. The entire logging process, after loading each core section onto the conveyor track, is controlled by the MST microprocessor with the display and recording of data being handled by the host computer.



**Figure 1**  
 Configuration of the Multi sensor core logger installed on the RV Marion Dufresne for the IMAGES MD101 North Atlantic Cruise.



## 1.2 Core handling.

Unsplit cores were delivered to the MST lab in 1.5 m sections. The cores temperature is low (few degrees Celsius) just after recovery but increases rapidly as soon as they are stored in the lab containers. Several hours (up to 24) of equilibration time proved necessary to bring the cores up to a stable room temperature before logging could begin. This equilibration is necessary to avoid that progressive temperature drifts take place during logging and affect physical properties results. At times, a temperature gradient of 4°C existed between the floor and the ceiling in the MST container. Because the container temperature could not be controlled accurately, it was necessary to store the cores in a horizontal position to ensure an homogeneous temperature along the length of the core.

## 1.3 Calibration and initial settings of the MST.

a) Core temperature - Changes in core temperature result in changes in P-wave velocities in the sediment which must be corrected for during processing. Sediment temperature was measured for each section and P-wave velocities were corrected using the formulae of Wilson (1960) and Wylie et al. (1956), and MacKenzie (1981).

b) External core diameter - Variations in core diameter cause variations in physical properties measurements which reflect the changing amount of sediment between the sensors instead of intrinsic sediment properties. For the P-wave velocity acquisition, the true diameter is measured for each measurement by an electronic caliper. For the gamma-ray attenuation measurements, the nominal external core diameter was reset after every four 1.5 m sections had been logged. Observed drift between calibrations was generally <0.2 mm. It is assumed that variations in external diameter are paralleled by changes in the internal diameter as it is not possible to measure this on a continuous basis. Only real changes in total offset are recorded.

c) Magnetic susceptibility - The drift of the susceptibility sensor over the course of the cruise was checked by measuring an iron ring of known susceptibility (527 cgs). The magnetic susceptibility of the core liners was measured by first setting the MS value to zero, then placing an empty section of liner inside the loop. During the cruise, two sets of PVC liner were used which didn't have the same physical properties signatures. Standards were constructed for both types of liner and different corrections were applied on bulk physical properties (i.e. the "old" liner appeared to have a higher magnetic susceptibility "background" level than the new one).

d) Gamma ray attenuation - Gamma ray attenuation was calibrated on a semi-weekly basis for the two liner types used (or whenever density and porosity values obtained for our water standard -which were run after every 4 sections- differed greatly from theoretically determined values). An empty section of core liner was placed between the source and the detector. Aluminium plates 6 mm thick (0, 1, then 2 up to 8) were placed inside the liner and the gamma ray attenuation measured. The resulting plot of density.vs.ln(gamma counts) provided the density calibration slope and intercept values used to correct GRAPE measurements for the attenuation of the liner.

Conversion of the gamma attenuation measurements to sediment density was done using the relationship established by Boyce (1976) for quartz.

The gamma ray beam produced on the shipboard MST could be collimated through one of two possible collimators (5 and 2.5 mm diameter). On this cruise only the larger diameter collimator was used due to the large diameter of the sediment cores.

## 1.4 MST Logging Procedure

The logging procedure used in the MST lab generally followed that outlined in the manufacturers instruction manual. The new GEOTEK system is designed to log an entire core at once by measuring on a continuous basis one section after the other. The major drawback of this method is that no data storage is performed on the harddrive of the host computer before the entire core is logged. Any computer problem arising during the logging procedure results, therefore, in the loss of the entire data set obtained so far. To avoid this problem, we scanned

the cores by intervals of four sections (a water standard was added on each 4 sections-run which allowed to have a close control on sensors calibration ). This procedure also enabled a close check to be kept of the cumulative error in core length caused by non-straight cuts of the liner (the core length was precisely reset at the beginning of each block of four sections). Sampling interval was set to **2 cm** for all cores.

The following parameters were used for most logging runs:

Maximum core length (for four sections plus water standard)	800 cm
Core sampling interval	2 cm
Nominal core diameter	113 mm
Core diameter logging	YES
Magnetic susceptibility units	SI
Magnetic susceptibility sampling time	1 Sec
A-Gamma count time	3 Sec

P-wave travel time offset 9.056 usec (varies)

Once a stable temperature had been reached, the unsplit sediment core was placed on the core track at the right hand end of the conveyor system, and aligned to its start position with one end of the core at a predetermined reference position. In this case a piece of empty core liner was used to separate the P-wave transducers and mark the reference position. The core then is moved incrementally along the length of the track, pushed by a block fixed to the conveyor belt, past or through the various sensors.

In order to ensure a good acoustic coupling between the P-wave transducers and the core, the liner was wiped with a wet sponge prior to logging and, when necessary, during logging by dripping a little distilled water onto the contact. The quality of the P-wave signal could be observed in "real time" on the oscilloscope integrated into the MST unit. As sections were stored in a horizontal position, there often proved to be air gaps between the sediment and the liner resulting in poor quality P-wave transmission.

Where possible, the endcaps were removed from the core sections to minimize errors in determining core length. Where this was not possible (for example, with fluid sediment, usually only in the top sections of the core), thin, temporary core caps were used (because the standard endcaps were too large to fit through the magnetic susceptibility loop) and their thickness noted. Occasionally core sections expanded out of the liner once the core caps were removed. In these cases, the amount of core extending beyond the liner was noted down.

Very badly damaged sections were not logged by the MST, the primary constraint being the ability of the section to fit through the magnetic susceptibility loop sensor (internal diameter 125 mm). Great care was taken when running damaged sections to ensure that the P-wave transducers were not "rammed" by following sections of core on the conveyor track.

## 1.5 Data processing

The MST generates three data files for each core logged; 1) a file containing the configuration setup for that particular core; 2) a "skewed" ASCII text file containing the raw data (explained below) and; 3) a random access data file generated as the core is being logged.

At this stage it was possible to correct logging errors by editing the text data file prior to processing. Usually logging errors were due to the temperature probe not being inserted in the core until after the section had begun to be logged. Occasional "double samples" caused by the core pusher jamming were also edited out at this stage.

As the MST's sensors are set at different positions along the conveyor track, every sample taken records values offset by a predetermined amount from the reference position. The MST itself only saves data in this "skewed" format and prior to any other processing, the data must first be unskewed. This was done using a processing package provided by the University of New Brunswick (written by Kate Jarrett). The same package was also used to calculate corrected

values for P-wave velocity, and Gamma attenuation for the effects of the liner and temperature variations. Density was also calculated as a function of gamma ray attenuation.

During processing a "Processing sheet" was kept on which all the relevant parameters were entered for each file. In general these consisted of:

- Water depth
- Depth of first sample
- Liner thickness
- Density intercept
- Density calibration slope
- P-wave travel time offset
- Bottom water temperature (assumed to be 4 C)
- Sediment density (assumed to be 2.7 g/cm<sup>3</sup>)
- Input filename
- Output filename

Processed data for each batch of four sections were saved with the water standard data included at the bottom of the file, as this provided a check on the accuracy of the calibrations being used. In addition a file containing data (with water values removed) for the entire core was accumulated by simply cutting and pasting the data for each group of four sections into one file. This file was then used to generate plots, using the Powermac application "Kaleidagraph".

## 1.6 Problems

No mechanical problems were experienced with the system during the cruise. On one occasion, a short circuit developed between the motor control unit and the conveyor track. This was fixed by simply cleaning and drying the connection.

Occasionally the core pusher proved insufficiently powerful to move two sections of core along the track. This problem was largely overcome by applying grease to the track to reduce friction.

Most problems were caused by software/computer errors, which disrupted the control system of the core logger. However, no computer related difficulties proved to be severe enough to prevent any core being logged.

The following notes refer mainly to the Microprocessor installed in the MST. This chip contains both RAM and PROM memory circuits. In order to operate the MST the instruction code for this microprocessor must first be downloaded from the host computer. Once installed in the PROM the code should then remain there without further downloading being necessary.

Initial software problems were an inability to access the test menu from the main menu. This was overcome by switching the system off (Computer and MST) and restarting. However, during the transfer of hex code between the host computer and MST when initializing the system, an error message suggesting that some of the code had become corrupted appeared (ERROR: BAD SYNTAX in line 9860).

A fault in the data display was also noted - sample number 13 was displayed twice on the screen every time it was recorded. Sample 13 itself was recorded without difficulty, however values for sample 14 were preceded by a repeat of those values recorded for sample 13.

As a result, only 12 cm of core could be logged before the system stopped. The motor would not advance and the software "froze", i.e. the data remained on the screen but nothing happened. Line 9860 in the MSCL21.I52 file on the C: drive was identical to that listed in the MST software printout. We also tried to start the MST from the A: drive with the same result (i.e. syntax error in the .I52 code). Therefore it appeared that the problem was with the software in the MST - ROM. An attempt to replace this with an MSCL21.I52 file off the backup disk (as per the instructions in the MST manual) was unsuccessful. We also tried to upload the MSCL21.I52 file from the ROM to compare its contents with those in the MST software manual and found that only the first half the code (up to line 5760, which is in mid-subroutine as well as being only halfway through the code) appeared to be present when the upload had been completed. An attempt to download uncorrupted code into the MST microprocessors RAM was

initially unsuccessful due to an apparent failure of the "New" command to reset the memory. This resulted in a MEMORY ALLOCATION ERROR, which prevented a large section of the code from being transferred.

## **2. Discrete measurements of index properties**

In order to calibrate correctly the continuous wet-bulk density records provided by the MST, discrete index properties measurements (bulk density, dry density, porosity) were performed on the IMAGES-MD101 cores (one measurement per section in most of the cores).

Calculations were based on the measurements of wet ( $W_m$ ) and dry ( $D_m$ ) mass of constant,  $10.36\text{-cm}^3$  plugs of sediment obtained with a precisely calibrated aluminium syringe. Dry weight were measured on samples after 24 hours drying in a  $60^\circ\text{C}$  oven. Sample mass was determined aboard the ship to a precision of  $\pm 0.01$  g using a Scitech electronic balance. This balance is equipped with a double-plateau. Acceleration changes due to the ship motion are instantaneously evaluated and corrected by measuring in the same time the weight of a known mass.

$$\text{Wet-bulk density (g/cm}^3\text{)} : W_m/10.36$$

$$\text{Dry-bulk density (g/cm}^3\text{)} : D_m/10.36$$

Porosity was calculated assuming a density of  $1.035$  g/cm<sup>3</sup> for pore water lost during drying. The volume of water was then calculated based on the weight loss : volume of pore water =  $W_m - D_m / 1.035$ . And thus, the porosity formula is:

$$\text{Porosity (\%)} : ((W_m - D_m) / 1.035) / 10.36 * 100$$

## **3. Spectrophotometry**

The Minolta CM-2002 hand-held spectrophotometer was used to measure reflected visible light on sediment cores that were splitted on board the Marion Dufresne during the cruise. The visible spectrum is divided in thirty one, 10-nm wide bands ranging from 400-700 nm. Measurements were usually performed every 5 cm all along the cores. The color reflectances were measured as soon as possible after the cores were split, because redox-associated color changes may occur rapidly when deep-sea sediments are exposed to the atmosphere.

A few problems were encountered at the beginning of the cruise and it became rapidly apparent that these problems resulted from an incorrect use of the Minolta which was not always pressed flush against the sediment surface during the measurements. Incorrectly measured sections were rapidly re-done after this problem has been evidenced by Dr. Ed Boyle who was in charge of the spectrophotometer acquisition.

The color reflectance profiles proved to be extremely powerful tools for preliminary stratigraphic and correlation purposes. Between  $40^\circ$  and  $55^\circ\text{N}$ , more particularly, color reflectance profiles have been successfully used for the direct tuning of the time scales by cyclostratigraphy in the precession and obliquity bands.

## **4. Video imaging and pictures**

In sediments from the North Atlantic, continuous records of color changes obtained by video proved to be extremely successful in picking rapid climatic variability (Cortijo et al., 1995). The system developed at the CFR was brought aboard the R/V Marion Dufresne for the IMAGES-MD101 cruise.

Working half sections were digitized by intervals of 75 cm in length with a tri-CCD (Charged Coupled Device) color camera. The lighting was constituted by two neon glow lamps. In the resulting video images, one pixel represents approximately 1.3 mm of sediment. Because of the

high sampling density of the video imaging, the resulting data files are huge and were stored on 1.2 Gigabytes magneto-optical disks.

Because of the time pressure during the cruise, the concatenation of the 75-cm images as well as the complete image processing could not be achieved on board the *Marion Dufresne*. The data processing will make it possible to eliminate illumination distortions which result in relatively strong, 75-cm cycles in the unprocessed video images.

We took color pictures of the 75-cm intervals that were scanned.

## **5. References**

- Boyce, R.E., 1976. Definitions and laboratory techniques of compressional sound velocity parameters and wet-water content, wet bulk density, and porosity parameters by gravimetric and gamma ray attenuation techniques. In Schlanger, S.O., Jackson, E.D., et al., *Init. Repts. DSDP*, 33: 931-958. Washington (U.S. Govt. Printing Office).
- Cortijo, E., Yiou, P., Labeyrie, L., and Cremer, M., 1995. Sedimentary record of rapid climatic variability in the North Atlantic Ocean during the last glacial cycle. *Paleoceanography*, 10(5), 911-926.
- MacKenzie, K. V., 1981. Nine-term equation for sound speed in the oceans, *J. Acoust. Soc. Am.*, 70(3), 807-812.
- Wilson, W. D., 1960. Speed of sound in sea water as a function of temperature, pressure and salinity, *J. Acoust. Soc. Am.*, 32(6), 641-644.
- Wylie, M. R. J., Gregory, A. R., and Gardner, L. W., 1956. Elastic wave velocities in heterogeneous and porous media, *Geophysics*, 21(1), 41-70.



## **B. Preliminary report on calcareous nannofossil biostratigraphy of IMAGES-MD101 cores**

Luc BEAUFORT and Jacques GIRAUDEAU

Calcareous nannofossil assemblages were examined in core catchers of cores retrieved during the IMAGES-MD101 cruise in order to provide age estimates for the core bases. Conventional biostratigraphy, using highest and lowest occurrence data of marker species, allows the identification of three biozones in the Late Pleistocene (NN21, NN20 and NN19; Martini, 1971). The NN19/20 boundary is marked by the last occurrence of *Pseudoemiliana lacunosa*, and the NN20/21 boundary corresponds to the first occurrence of *Emiliana huxleyi*. In order to increase the stratigraphic resolution, it is possible to use the "acmes" of some coccolithophore species.

Works of Pujos (1988) and Weaver (1993), give a stratigraphic framework based on isotopic stages that we used during the MD-IMAGE 101 cruise:

- 1 acme of *Emiliana huxleyi* during Stage 1 to 3
- 2 acme of *Gephyrocapsa mulleare* during Stage 5
- 3 acme of small *Gephyrocapsa* spp during Stage 6 to 8
- 4 acme of *Gephyrocapsa caribbeanica* during Stage 9 to 15

In addition to this framework, a higher resolution can be achieved by looking at the abundances of coccoliths which proved to be usually higher during odd isotopic stages in the North Atlantic Ocean.

The following table summarizes our observations using conventional biostratigraphy, acmes and abundance of coccolithophores.

Note : If conventional biostratigraphy (LAD, FAD) provides a reliable biochronological scale (the LAD of *P. lacunosa* and the FAD of *E. huxleyi* have been recognized as widely spread isochronous datums), the use of acmes and abundances is a recent improvement of the coccolith biostratigraphy. Their dependence on local ecological factors which could result in chronological discrepancies, has not been tested enough yet.

Martini, E., Standard Tertiary and Quaternary calcareous nannoplankton zonation, In *Proceedings of the 2nd plankton conference, Roma*, A. Farinacci (ed), pp. 739-777, Roma, 1971.

Pujos, A., Spatio-temporal distribution of some Quaternary coccoliths, *Oceanolog. Acta*, 11, 65-76, 1988.

Weaver, P.P.E., High resolution stratigraphy of marine Quaternary sequences, In *High resolution stratigraphy*, 70, E.A. Hailwood and R.B. Kidd (ed), pp. 137-153, Geophysical Society Special Publication, 1993.

CORE (sample)	Depth	Martini Zones	Coccoliths	Iso. stage
MD95 2001 (cc)	22.37	NN21	>G.m.; E.h.	5
MD95 2002 (2890)	29.99	NN21	>G.m.; E.h.; RW	5
MD952003 (cc)	27.55	NN19	low abundance	6
MD95 2004 (cc)	28.48	NN20	> G.c.	13
MD95 2005 (cc)	22.02		barren	9 (or 10-11)
MD95 2008 (cc)	21.50		rare- small G.	
MD95 2009 (cc)	32.52		barren	
MD95 2010 (cc)	17.65	NN20	>G.c.; small G.	8-10
MD95 2011 (cc)	19.77	NN21	>G.m.; E.h.; RW	3-4
MD95 2012 (cc)	13.10	NN21	>RW (>G.m.; E.h.)	
MD95 2013 (cc)	33.54	NN21	>G.m. E.h.	5
MD95 2014 (3356)	34.50	NN19	>G.c.; ?P.L.	13
MD95 2015 (3267)	32.86	NN19	>G.c.; ?P.L.	13
MD95 2016 (3190)	32.86	NN19	>G.c.; ?P.L.	13
MD95 2017 (cc)	34.58	NN18	>G.c.(abund.)	9 or 11
MD95 2019 (1210)	12.86		barren	
MD95 2020 (cc)	8.63		barren	
MD95 2021 (cc)	4.48	NN21	>E.h., low abund.	2-3
MD95 2022 (cc)	18.34	NN19	>G.c.; P.L.	13 or 15
MD95 2023 (cc)	34.68		barren	
MD95 2027 (cc)	46.70	NN19	>G.c.; P.L.	13
MD95 2030 (cc)	35.56	NN19	P.L.; Low abund,RW	>12
MD95 2030 (cc)	27.00		RW	
MD95 2031 (cc)	27.00			
MD95 2034 (cc)	47.16	NN21	small G, L.ab.	6
MD95 2035 (cc)	39.25	NN20	>G.c.	9
MD95 2036 (cc)	52	NN21	>small G.abund.	7
MD95 2037 (cc)	35.81	NN19	>G.c; P.L.	13-15
MD95 2038 (cc)	37.86	NN19	>G.c; P.L.	13-15
MD95 2039 (cc)	35.73	NN20	>G.c.(low abund.)	10
MD95 2040 (cc)	35.24	NN20	>G.c. (abund.)	9 or 11
MD95 2041 (cc)	17.62	NN21	>G.m.	5
MD95 2042 (cc)	39.58	NN21	>small G.	7
MD95 2043 (cc)	37.40	NN20	>sm.G.+ G.c+RW	8

Table 1: Coccolith biostratigraphy for the base of the cores retrieved during MD-IMAGE 101 cruise: > = acme; small G. = small *Gephyrocapsa* sp., G.c. = *G.caribbeanca*; G.m. = *G.muelleriae*; E.h. = *E. huxleyi*; P.L. = *Pseudoemiliania lacunosa*; RW = reworked species (pre-Pleistocene); HO = highest occurrence, L.ab. = low abundance.





## SITE REPORTS

The report for each site contains the following pieces of information:

**\* First page :**

- a site description (location, water depth, length of core);
- a detailed map showing the core(s) location;
- 3.5 kHz site survey (when available).

**\* Second page :**

- selected, high-resolution physical properties records
  - magnetic susceptibility;
  - GRAPE density;
  - color reflectance - parameter a, b or L




**\* Third page :**

- the entire set of physical properties records obtained on board
  - magn. suscepti., acoustic veloc., GRAPE density (from MST);
  - L, a and b parameters (color reflectance data)

**\* Fourth page :**


- sedimentological description of the core


Core description

<p><b>Constituents</b></p> <p>Nanofossils </p> <p>Siliceous skeletons </p> <p>Detrital carbonate grains ("detrital carbonate") </p> <p>carbonated silty clay (biogenic and detrital carbonate) </p> <p>abundant volcanic glass </p> <p>Dropstone </p> <p><i>The sediments were qualified according to the dominant constituents and the texture of the lithic fraction. Clayey ooze : &gt;50 % biogenic skeletons, clay dominates the lithic fraction.</i></p>	<p><b>Structure</b></p> <p>burrowing </p> <p>Iron sulfide (FeS) dots and patches concentrated in indistinct layering </p> <p>distinct-indistinct "Greenish layers" </p> <p>distinct-indistinct lamination </p>
<p><b>Texture</b></p> <p>Clay </p> <p>Silty clay </p> <p>clayey silt </p> <p>silt </p> <p>Sand </p> <p>mud (clay+silt+sand) </p>	<p><b>Coring disturbances</b></p> <p>Slightly disturbed </p> <p>Soupy </p> <p>highly disturbed </p> <p>flow in </p> <p>void </p>
<p><b>Colour</b></p> <p>SY4/2 Munsell colour chart reference</p> <p>Sharp change </p> <p>Gradual change </p>	

le 09/07/95

MD101 IMAGES - Mai/Juillet 95







lat: 46°48.17 N - long: 08°40.28 W  
 water depth: 3788 m  
 length: 22.37 m

# Core MD 95-2001


0 20 40 60 80 100 1,4 1,6 1,8 2 2,2 -2 -1 0 1 2 3



MD 95-2001



MD 95-2001




# Core MD 95-2002


lat: 47°27.12 N - long: 08°32.03 W

water depth: 2174 m


length: 29.99 m

IMAGES-MD101






MD 95-2002





MD 95-2002




lat: 55°10.61 N - long: 14°45.32 W


water depth: 2365 m  
length: 26.93 m

# Core MD 95-2003






MD 95-2003




MD 95-2003




# Core MD 95-2004

lat: 55°28.07 N - long: 14°40.51 W  
water depth: 2177 m  
length: 28.485 m




MD 95-2004




# MD95-2004

60

IMAGES-MD101



MD 95-2004



# Core MD 95-2005




62

lat: 57°02' N - long: 10°03.83' W




water depth: 2130 m

length: 22.02 m


IMAGES-MD101




0 50 100 150 200 1,6 1,8 2 2,2 2,4 -0,5 0 0,5 1 1,5 2



MD 95-2005





MD 95-2005






# Core MD 95-2006


lat: 57°01.81 N - long: 10°03.48 W  
water depth: 2122 m  
length: 30.22 m



0 100 200 300 1,6 1,8 2 2,2 2,4 1450 1475 1500 1525 1550



MD 95-2006



**Core MD95-2006 not opened on board.**

**No sediment description.**


# Core MD95-2007

IMAGES-MD101

lat: 57°31.06 N - long: 08°23.17 W

water depth: 158 m

length: 19.35 m



**Core MD95-2007 not measured on MST.**



**Core MD95-2007 not opened on board.**

**No core description.**



# Core MD 95-2008

IMAGES-MD101




lat: 62°44.35 N - long: 03°59.64 W

water depth: 1016 m


length: 21.5 m





0 300 600 900 1200 1,4 1,6 1,8 2 2,2 -1 -0,5 0 0,5 1



MD 95-2008




MD 95-2008






# Core MD 95-2009

IMAGES-MD101


lat: 62°44.25 N - long: 03°59.86 W  
water depth: 1027 m  
length: 32.43 m




0 500 1000 1500 1,4 1,6 1,8 2 2,2 -1 -0,5 0 0,5 1



MD 95-2009





MD 95-2009




# Core MD 95-2010


IMAGES-MD101

lat: 66°41.05 N - long: 04°33.97 E  
Water depth: 1226 m  
length: 32.53 m



0 50 100 150 200 250 1,4 1,6 1,8 2 2,2 -2 -1 0 1 2









# Core MD 95-2011

IMAGES-MD101




lat: 66°58.18 N - long: 07°38.36 E

water depth: 1048 m


length: 17.49 m





0 40 80 120 160 1,4 1,6 1,8 2 2,2 2,4 -1 0 1



MD 95-2011



MD 95-2011







a

Dens.


IMAGES-MD101  
Suscept.






MD 95-2012

MD 95-2012  
Section IX to XV



MD 95-2012  
Section I to VIII




# Core MD 95-2013

lat: 72°10.917 N - long: 13°05.632 E

water depth: 1638 m


length: 13.1 m



**Core MD95-2013 had a badly damaged liner, it was not measured on the MST.**



MD 95-2013





# Core MD 95-2014

IMAGES-MD101




lat: 60°34.93 N - long: 22°04.52 W


water depth: 2397 m

length: 33.61 m




0 250 500 750 1,4 1,6 1,8 2 30 35 40 45 50 55






MD 95-2014


Section XVI-XXIII



Section IX to XV



Section I to VIII






# Core MD 95-2015

IMAGES-MD101

lat: 58°45.74 N - long: 25°57.54 W

water depth: 2630 m




length: 34.42 m




0 100 200 300 400

1,2 1,4 1,6 1,8


40 50 60



MD 95-2015



MD 95-2015





# Core MD 95-2016


IMAGES-MD101


lat: 57°42.46 N - long: 29°25.44 W

water depth: 2318 m


length: 32.83 m







MD 95-2016




# Core MD 95-2017


IMAGES-MD101


lat: 53°02.56 N - long: 33°31.51 W

water depth: 3100 m

length: 34.59 m







**Core MD95-2017 not opened on board.**


**No sediment description.**

# Core MD 95-2018

lat: 50°43.88 N - long: 43°08.54 W

water depth: 4221 m




length: 14.7 m




0 100 200 300 400 500

1,6 1,8 2


35 40 45 50 55



MD 95-2018



MD 95-2018



Detrital carbonate  
clayey silt

Sediment in bags

527-554  
Slabs 554-581.5  
581-600

Detrital carbonate rich mud-  
**turbidites**  
clayey silt

Sandy silty clay 683-710  
Slabs 710-738  
738-750  
gneiss Ø 6 cm


No description of sections VII, VIII, IX, X  
Core base: 1470 cm

# Core MD 95-2019

lat: 51°05.68 N - long: 3°13.27 W

water depth: 4262 m




length: 12.86 m




0 100 200 300 400 500

1,6 2 2,4

30 35 40 45 50 55




MD 95-2019



MD 95-2019

Section I to VIII




# Core MD 95-2020

lat: 51°02.42 N - long: 43°00.62 W

water depth: 4289 m

length: 8.63 m






Suscept.


Dens.

L


0 200 400 600 800 1,6 1,8 2 2,2 2,4 2,6 30 35 40 45 50



MD 95-2020






MD 95-2020







0 100 200 300 400 500 1,4 1,6 1,8 2 2,2 35 40 45 50 55



MD 95-2021



MD 95-2021






# Core MD 95-2022


lat: 50°34.38 N - long: 43°03.51 W


water depth: 4245 m

length: 18.37 m




0 200 400 600 1,5 1,75 2 2,25 2,5 30 40 50 60






MD 95-2022

Section IX to XVI



Section I to VIII





# Core MD 95-2023

lat: 50°58.32 N - long: 43°13.04 W

water depth: 4198 m


length: 34.68 m




L

Dens.

IMAGES-MD101  
Suscept.








# Core MD 95-2024


IMAGES-MD101


lat: 50°12.40 N - long: 45°41.22 W

water depth: 3539 m

length: 29.52 m







**Core MD95-2024 not opened on board.**



**No sediment description.**

# Core MD 95-2025

lat: 49°47.65 N - long: 46°41.85 W

water depth: 3009 m

length: 35.12 m





L

Dens.

IMAGES-MD101  
Suscept.

0 200 400 600 1,5 2 2,5






h

**Core MD95-2025 not opened on board.**

**No sediment description.**

# Core MD 95-2026


lat: 48°13.62 N - long: 46°39.39 W  
water depth: 878 m  
length: 27.91 m




L

Dens.

IMAGES-MD101  
Suscept.





**Core MD95-2026 not opened on board.**

**No sediment description.**


# Core MD 95-2027

IMAGES-MD101

lat: 41°44.67 N - long: 47°24.79 W

water depth: 4112 m




length: 47.67 m




0 50 100 150 200

1,6 1,8 2

40 45 50 55 60










# Core MD 95-2028

lat: 42°05.99 N - long: 52°44.85 W

water depth: 3368 m

length: 34.2 m




L


Dens.

IMAGES-MD101  
Suscept.

0 50 100 150 200 1,5 1,75 2 2,25



MD 95-2028



**Core MD95-2028 not opened on board.**

**No sediment description.**




# Core MD 95-2029

IMAGES-MD101


lat: 43°06.78 N - long: 55°15.96 W

water depth: 4156 m

length: 35.1 m



0 50 100 150 1 1,2 1,4 1,6 1,8 2





**Core MD95-2029 not opened on board.**

**No sediment description.**



# Core MD 95-2030

IMAGES-MD101

lat: 44°19.14 N - long: 53°43.44 W

water depth: 1573 m


length: 35.59 m




**Core MD95-2030 not measured on the MST.**



Section I to VIII




Section IX to XVII



Section XVIII to XXII

MD 95-2030




# Core MD 95-2031

IMAGES-MD101

lat: 44°18.45 N - long: 53°44.14 W

water depth: 1570 m


length: 27.72 m




L

Dens.

IMAGES-MD101  
Suscept.



MD 95-2031



**Core MD95-2031 not opened on board.**




**No sediment description.**

# Core MD 95-2032

lat: 44°39.46 N - long: 55°37.14 W

water depth: 1433 m

length: 28.54 m



**Core MD95-2032 not measured on MST.**



**Core MD95-2032 not opened on board.**




**No sediment description.**

# Core MD 95-2033

lat: 44°39.87 N - long: 55°37.21 W



water depth: 1412 m


length: 29.68 m



0 45 90 135 180

1,6 2 2,4





**Core MD95-2033 not opened on board.**



**No sediment description.**

# Core MD 95-2034

lat: 33°41.46 N - long: 57°34.55 W

water depth: 4461.5 m


length: 47.16 m




L


Dens.

IMAGES-MD101  
Suscept.





MD 95-2034





# Core MD 95-2035


lat: 33°29.20 N - long: 57°53.16 W

water depth: 4286 m


length: 39.25 m







MD 95-2035







# Core MD 95-2036

lat: 33°41.44 N - long: 57°34.55 W


water depth: 4461.5 m

length: 52.64 m









# MD95-2036



MD 95-2036

Unopened sections I to XXVI: 0-4050 cm  
and XXXI to XXXVI: 4500-5264 cm






# Core MD 95-2037

IMAGES-MD101

lat: 37°05.23 N - long: 32°01.87 W

water depth: 2159 m

length: 35.81 m




L


Dens.

IMAGES-MD101  
Suscept.


0 25 50 75 100 1,2 1,4 1,6 1,8 2



MD 95-2037




MD 95-2037






# Core MD 95-2038

IMAGES-MD101


lat: 37°45.15 N - long: 27°11.21 W  
water depth: 2310 m  
length: 37.85 m




0 100 200 300 400 500 1,4 1,6 1,8 2 20 40 60 80



MD 95-2038



MD 95-2038





# Core MD 95-2039




lat: 40°34.71 N - long: 10°20.91 W

water depth: 3381 m


length: 35.71 m




0 20 40 60 80 1,6 1,8 2 2,2 30 40 50 60 70



MD 95-2039



MD 95-2039




**Clayey nannofossil ooze**  
- with detrital carbonates

- black spots

**Nanno-bearing silty clay**

- FeS spots scattered throughout  
- homogenous  
- some sponges spicules




- long vertical burrow filled by fine grained pyrite

**Nanno-bearing silty clay**

**silty clay**  
sand layer, mainly quartz grains and detrital carbonates

**Nanno-bearing silty clay**



**Interbedded Nanno-bearing silty clay and clayey Nanno ooze as shown by color changes**

- with detrital carbonates

- numerous black spots

- several green layers

- numerous green layers

- thin fine grained sand layer (5 Y 4/2)

Stiff

- sandy

- numerous black spots

- silty clay (5 Y 4/2)

3573 cm

# Core MD 95-2040


202



IMAGES-MD101

lat: 40°34.91 N - long: 09°51.67 W


water depth: 2465 m

length: 35.24 m






MD 95-2040



MD 95-2040




# Core MD 95-2041

IMAGES-MD101

lat: 37°50 N - long: 09°30.64 W

water depth: 1123 m




length: 17.62 m




0 50 100 150 200

1,4 1,6 1,8 2 2,2 2,4


40 42,5 45 47,5 50



MD 95-2041



MD 95-2041



1762 cm



# Core MD 95-2042

IMAGES-MD101


lat: 37°47.99 N - long: 10°09.99 W


water depth: 3146 m

length: 39.56 m



0 50 100 150 200 250 1,5 1,6 1,7 1,8 1,9 2 2,1





**Core MD95-2042 not opened on board**



**No sediment description.**

# Core MD 95-2043



lat: 36°08.60 N - long: 02°37.27 W

water depth: 1841 m

length: 37.46 m



0 10 20 30 40 50 60 1,4 1,6 1,8 2 2,2





**Core MD95-2043 not opened on board.**

**No sediment description.**

